

# Optimal care pathway for people with biliary cancer

FIRST EDITION





# Optimal care pathway for people with biliary cancer

FIRST EDITION

Endorsed by



Australian Government  
Cancer Australia



## Statement of acknowledgement

We acknowledge the Traditional Owners of Country throughout Australia and their continuing connection to the land, sea and community. We pay our respects to them and their cultures and to Elders past, present and emerging.

This work is available from the Cancer Council website <<https://www.cancer.org.au/health-professionals/optimal-cancer-care-pathways>>.

First edition published in March 2026.

ISBN: 978-1-7644010-6-7

Pancare Foundation 2025, *Optimal care pathway for people with biliary cancer, 1st ed.*, Pancare, Heidelberg.

Enquiries about this publication can be sent to <[Melanie.Regan@pancare.org.au](mailto:Melanie.Regan@pancare.org.au)> or <[OCPs@canceraustralia.gov.au](mailto:OCPs@canceraustralia.gov.au)>.

# Contents

## Contents

<b>Welcome and introduction</b>	<b>i</b>
<b>Summary</b>	<b>1</b>
<b>Intent of the optimal care pathways</b>	<b>7</b>
Optimal care pathway resources	8
<b>Principles of the optimal care pathway</b>	<b>10</b>
Principle 1: Patient-centred care	10
Principle 2: Safe and quality care	12
Principle 3: Multidisciplinary care	13
Principle 4: Supportive care	14
Principle 5: Care coordination	15
Principle 6: Communication	16
Principle 7: Research and clinical trials	18
<b>Summary – optimal timeframes</b>	<b>19</b>
<b>Optimal care pathway</b>	<b>20</b>
Step 1: Prevention and early detection	21
Step 2: Presentation, initial investigations and referral	22
Step 3: Diagnosis, staging and treatment planning	25
Step 4: Treatment	34
Step 5: Care after initial treatment and recovery	45
Step 6: Managing recurrent, residual or metastatic disease	51
Step 7: End-of-life care	55
<b>Contributors and reviewers</b>	<b>58</b>
<b>Appendix A: Supportive care domains</b>	<b>60</b>
<b>Appendix B: Psychological needs</b>	<b>61</b>
<b>Appendix C: Special population groups</b>	<b>62</b>
<b>Appendix D: Complementary therapies</b>	<b>69</b>
<b>Appendix E: Members of the multidisciplinary team for biliary cancer</b>	<b>70</b>
<b>Resource list</b>	<b>71</b>
<b>Glossary</b>	<b>75</b>
<b>References</b>	<b>77</b>

# Welcome and introduction

On behalf of the Biliary Cancer Expert Working Group and the larger Biliary Tract Cancer Consensus Guidelines Group, we are proud to present the first edition of the optimal care pathway for people with biliary cancer.

Some cancers are simpler to treat than others, but all are complex. However, the principles of high-quality care are similar for all cancers and, if followed, are highly likely to achieve the best possible outcomes for patients and their families and carers.

Despite Australia having a world-leading health system, navigating through it can be complicated. Care is provided across the country by large and small teams and services, and given resources are limited, it is inevitable that variation exists. The optimal care pathways for cancer care allow the entire community to understand what good-quality care looks like and to put in place systems and processes that help ensure this is achieved.

This complexity in care has led us to develop the optimal care pathways. These pathways describe an integrated model of care incorporating both the technical aspects of care and the delivery of such care. They provide a national standard for the high-quality cancer care that all Australians should expect.

The optimal care pathways are not intended to replace clinical guidelines. The choice of treatment for an individual is a professional responsibility based on patient factors and preferences, practice guidelines and emerging data. But the optimal care pathway for biliary cancer will help ensure improved cancer care and outcomes for people with biliary cancer.

The optimal care pathways should be read and understood by all health professionals involved in cancer care, as well as trainees in all relevant disciplines. We also recommend them to patients and their carers in the hope that they may guide discussions with their healthcare team. There are specific versions of the optimal care pathways for consumers called the guides to best cancer care. These are available in different languages.

This optimal care pathway was formally endorsed by the Cancer and Population Screening Committee following advice from Cancer Australia's National Cancer Expert Group. Formal committee endorsement acknowledges the importance of this optimal care pathway and the responsibility of the health system to deliver care in an appropriate, culturally safe, responsive and coordinated way that ensures equal and timely access and that all Australians experience optimal health outcomes.

We would like to reiterate our thanks to everyone involved for generously dedicating considerable time and effort in developing this optimal care pathway. This includes many people and organisations that contributed to reviewing these guides and the strong support of the federal, state and territory governments. It is our sincere hope that it will point to the importance of patient-centred cancer care and consequently help smooth the difficult road that patients and their carers must walk with a cancer diagnosis.



**Professor John Zalcberg AO**

Chair, Biliary Cancer Expert Working Group

Co-chair, Biliary Tract Cancer Consensus Guidelines Group

# Summary

Support: Assess supportive care needs at every step of the pathway and refer to appropriate health professionals or organisations.

The optimal care pathways describe the standard of care that should be available to all cancer patients treated in Australia. The pathways support patients and carers, health systems, health professionals and services, and encourage consistent optimal treatment and supportive care at each stage of a patient's journey. Seven key principles underpin the guidance provided in the pathways: patient-centred care; safe and quality care; multidisciplinary care; supportive care; care coordination; communication; and research and clinical trials.

This quick reference guide provides a summary of the *Optimal care pathway for people with biliary cancer*. Biliary cancer includes cancer of the gallbladder and biliary tract cancer (also called cholangiocarcinoma).

**Please note that not all patients will follow every step of the pathway.**

## Step 1: Prevention and early detection

### Prevention

The causes of biliary cancer are not fully understood, and there is currently no clear prevention strategy.

### Risk factors

There are very few people at high risk. People with the following risk factors are at increased risk:

#### Gallbladder cancer

- Gallstones
- Having gallbladder polyps
- History of long-term inflammation of the gallbladder
- First-degree or immediate relative (mother, father, sibling or child) having gallbladder cancer
- Inherited genetic conditions such as abnormalities of the liver, gallbladder or bile ducts
- Obesity
- Smoking
- Being from East Asia or South America.

#### Biliary tract cancer

- Medical history of long-term inflammation of the bile duct such as primary sclerosing cholangitis
- Biliary cysts
- Inherited genetic conditions such as abnormalities of the liver, gallbladder or bile ducts
- Chronic liver disease such as cirrhosis, hepatitis B infection and hepatitis C infection
- Infection with a liver fluke parasite (linked to bile duct cancer cases in Africa and Asia)

- Genetic changes such as hereditary nonpolyposis colorectal cancer syndrome (Lynch syndrome), characterised by abnormalities of the mismatch repair genes (MMR) *MLH1*, *MSH2*, *MSH6* and *PMS2*, as well as mutations in *BRCA1/2*
- Exposure to Thorotrast, which is a radioactive compound that was used in x-rays from the 1930s to the 1950s
- Exposure to asbestos
- Being from East Asia or South America
- Pre-existing inflammatory bowel disease such as chronic ulcerative colitis
- Older age
- Diabetes
- Obesity.

### Early detection

Being aware of the risk factors and who is at high risk can guide monitoring and referral to a specialist.

### Screening recommendations

Routine screening for biliary cancer is not currently recommended in either the general population or in relatives of people with biliary cancer.

For people at higher risk, such as those with primary sclerosing cholangitis, long-term inflammation of the gallbladder or gallbladder polyps (6–10 mm), consider annual surveillance using ultrasound. Consider cholecystectomy for patients with gallbladder polyps over 10 mm.

### Checklist

- Current and past medical history discussed, including any genetic and inherited conditions.
- Recent weight changes discussed, and the patient's weight recorded.
- Alcohol intake discussed and recorded and support for reducing alcohol consumption offered if appropriate.
- Smoking status discussed and recorded, and brief smoking cessation advice offered to smokers.
- Physical activity recorded.
- Referral to a dietitian considered.
- Referral to a physiotherapist or exercise physiologist considered.
- Education on being sun smart considered.
- Any recent travel to Asia or Africa where liver fluke exposure may occur, discussed.
- Potential exposure to Thorotrast and asbestos noted.

## Step 2: Presentation, initial investigations and referral

Many cases present with non-specific symptoms or are asymptomatic until advanced stages of the disease.

The following signs and symptoms should be investigated:

- jaundice (requires urgent investigation) with or without darkened urine, pale bowel movements and itchy skin
- unexplained nausea or vomiting
- unexplained weakness
- unexplained loss of appetite and weight loss
- fevers and chills
- pain in the right side of the abdomen.

### Initial investigations

The patient's GP should take a thorough medical history to check for any risk factors and symptoms that suggest biliary cancer.

Where there is a suspicion of biliary cancer, consider:

- abdominal CT scan and/or abdominal ultrasound
- serum CA 19-9, liver function tests, full blood examination, biochemistry and coagulation screen (INR/APTT).

Where jaundice is present, the following should be performed, reviewed and the patient referred to a specialist doctor urgently:

- liver function tests and coagulation screen (INR/APTT)
- abdominal ultrasound
- CT of the abdomen and pelvis including a multiphase scan of the liver, provided this does not delay urgent referral.

### Referral options

At the referral stage, the patient's GP or other referring doctor should advise the patient about their options for referral, waiting periods, expertise, if there are likely to be out-of-pocket costs and the range of services available. This will enable patients to make an informed choice of specialist and health service.

### Communication

#### The GP's responsibilities include:

- explaining to the patient and/or carer who they are being referred to and why
- supporting the patient and/or carer while waiting for specialist appointments
- informing the patient and/or carer that they can contact the Cancer Council on 13 11 20 and/or the Pancare Foundation on 1300 881 698 and/or the Liver Foundation on 1800 841 118.

### Checklist

- Signs and symptoms recorded.
- Patient urgently referred to a biliary cancer specialist.
- Supportive care needs assessment completed and recorded, and referrals to allied health services actioned as required.
- Patient notified of support services such as the Cancer Council 13 11 20 and/or the Pancare Foundation on 1300 881 698 and/or the Liver Foundation on 1800 841 118.
- Referral options discussed with the patient and/or carer including cost implications.

### Timeframe

Patients who present with jaundice should be referred for tests immediately and followed up rapidly.

Other symptoms require investigations to be completed and reviewed by a GP **within two weeks** of the first appointment.

Patients with suspected or proven biliary cancer should be immediately referred and seen by a gastroenterologist, hepatologist, oncologist or hepatopancreatobiliary (HPB) surgeon with expertise in biliary cancer management and linked to a multidisciplinary team **within two weeks** of referral to the specialist.

## Step 3: Diagnosis, staging and treatment planning

### Diagnosis and staging

The minimum diagnostic procedure that should be completed before the multidisciplinary meeting (MDM) is a CT scan. The other tests can be performed under the direction of the MDM and should not delay presentation of a patient with suspected biliary cancer to the MDM. If the remaining staging tests reveal contrary information the patient may be re-presented to the MDM for further consideration.

Diagnosis of a mass is primarily by imaging. Contrary to the situation with most cancers, definitive tissue diagnosis before multidisciplinary planning is not recommended. Biopsy approaches may be complex but are generally required where there is diagnostic uncertainty, or where tissue is required for systemic therapy (such as chemotherapy) or clinical trials. This should be facilitated and overseen by the MDM process. Final pathology may, rarely, prove benign, despite appropriate work-up.

Contrast-enhanced CT scan of the chest, abdomen and pelvis, including multiphase CT scan of the liver is recommended (to be completed first if not already performed).

The following investigations are suggested and can occur under the direction of the MDM:

- contrast-enhanced MRI liver and magnetic resonance cholangiopancreatography (MRCP)
- endoscopic ultrasound (EUS)
- FDG-PET. (This can be useful in some cases but is not necessary in all cases. It may be a useful tool, for example, to evaluate evidence of spread of intrahepatic tumours or equivocal findings on radiology.)

Urgent stenting is **not** recommended if the patient's symptoms are mild. An appropriate diagnostic work-up should take place first. Stenting should not be done before the patient is reviewed at an MDM to determine if the tumour is resectable.

Staging for biliary cancer is vital to determine if the disease is resectable or non-resectable because the treatment pathways are different and preparation for resectable disease is highly specific.

A laparoscopy may be included in staging for resectable biliary cancer. Obtain a core biopsy for diagnostic pathology and molecular profiling before starting systemic or locoregional therapies.

### Genetic testing

Inherited genetic conditions causing biliary cancer are uncommon.

The types of genetic changes linked with an increased risk of biliary cancer include Lynch syndrome, characterised by abnormalities of the mismatch repair genes (MMR) *MLH1*, *MSH2*, *MSH6* and *PMS2*, as well as mutations in *BRCA1/2*.

Patients with a strong family history of cancer should be referred to a familial cancer clinic to determine whether the tumour is part of an inherited predisposition to cancer.

### Treatment planning

Patients with proven or suspected biliary cancer should be discussed at an MDM **within two weeks** of the initial specialist review. The MDM can occur after the minimum diagnostic tests; the full range of tests should not delay presentation at an MDM.

### Research and clinical trials

Consider enrolling the patient in clinical trials where available and appropriate. Search for a trial <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>.

### Communication

#### The lead clinician's<sup>1</sup> responsibilities include:

- discussing a timeframe for diagnosis and treatment options with the patient and/or carer
- explaining the role of the multidisciplinary team in treatment planning and ongoing care
- encouraging discussion about the diagnosis, prognosis, advance care planning and palliative care while clarifying the patient's wishes, needs, beliefs and expectations, and their ability to comprehend the communication
- providing appropriate information and referral to support services as required
- communicating with the patient's GP about the diagnosis, treatment plan and recommendations from MDMs.

### Checklist

- Diagnosis confirmed.
- Performance status and comorbidities measured and recorded.
- Patient discussed at an MDM and decisions provided to the patient and/or carer.
- Clinical trial enrolment considered.
- Supportive care needs assessment completed and recorded and referrals to allied health services actioned as required.
- Patient referred to support services (Cancer Council, Pancare Foundation, Liver Foundation) as required.
- Treatment costs discussed with the patient and/or carer.

### Timeframe

Diagnostic and staging investigations that the MDM has recommended should be completed **within two weeks** of referral.

<sup>1</sup> Lead clinician – the clinician who is responsible for managing patient care. The lead clinician may change over time depending on the stage of the care pathway and where care is being provided.

## Step 4: Treatment

### Establish intent of treatment

- Curative
- Anti-cancer therapy to improve quality of life and/or longevity without expectation of cure
- Symptom palliation

A treatment plan may also be informed by genomic testing. Genomics-informed cancer care can improve the experience of people affected by cancer by optimising the treatment response and minimising treatment toxicity.

### Treatment options

#### Surgery

- For resectable disease surgical resection offers the best long-term survival chance in patients with biliary cancer. Palliative surgery for metastatic cancer is not usually recommended.

#### Intrahepatic biliary cancer

- Partial hepatectomy (liver resection) is recommended.

#### Perihilar biliary cancers

- Right extended hepatectomy or left hemi-hepatectomy is recommended.

#### Distal biliary cancer

- Pancreaticoduodenectomy (Whipple's procedure) is recommended.

#### Gallbladder cancer

- Cholecystectomy including removing surrounding tissue such as the lymph nodes, adjacent bile ducts and part of the liver is recommended.

### Liver transplantation

- In the case of intrahepatic biliary cancer, consider a liver transplant for lesions smaller than 2 cm.
- For perihilar biliary cancer, consider a liver transplant for lesions no larger than 3 cm.
- For both sites of biliary cancer, there needs to be clear evidence that there are no metastasis including lymph node metastasis before considering a liver transplant.

### Locoregional therapy

#### Radiation therapy

- External beam radiation therapy (EBRT), chemoradiation (CRT) or stereotactic body radiotherapy / stereotactic ablative radiotherapy (SBRT/SABR) may be appropriate.
- Radiotherapy may be considered for patients with:
  - unresectable/locally advanced biliary cancer after completing systemic therapy and the disease is stable
  - intrahepatic biliary cancer with or without vascular involvement for patients in whom surgery is contraindicated
  - metastatic intrahepatic biliary cancer to reduce the risk of liver failure from local disease progression.
- Adjuvant CRT may be considered after margin positive resection (R1) or resection of node-positive disease.
- Adjuvant CRT may be considered following R0 resection of T2 or N1 gallbladder cancer.

#### Intra-arterial treatment

- May be either transarterial chemoembolisation (TACE), selective internal radiation therapy (SIRT) or hepatic arterial infusion chemotherapy (HAIC).

#### Thermal ablation

- Radiofrequency ablation (RFA) or microwave ablation (MWA) may be appropriate.
- Consider this for patients with intrahepatic biliary cancer no greater than 3 cm who have contraindications to surgery.

### Checklist

- Intent of treatment established.
- Risks and benefits of treatments discussed with the patient and/or carer.
- Treatment plan discussed with the patient and/or carer.
- Treatment plan provided to the patient's GP.
- Treating specialist has adequate qualifications, experience and expertise.
- Supportive care needs assessment completed and recorded, and referrals to allied health services actioned as required.
- Early referral to palliative care considered.
- Advance care planning discussed with the patient and/or carer.

## Step 4: Treatment continued

### Systemic therapy

- Adjuvant chemotherapy is recommended for all patients following resection for biliary cancer.
- Palliative chemotherapy is recommended for most patients with advanced biliary cancer, unless contraindicated by frailty or comorbidities.
- There are several approved targeted therapies available to treat patients with locally advanced, unresectable or metastatic biliary cancer, with more undergoing clinical trials. Molecular profiling is advised for all patients to guide appropriate targeted and immunotherapy choice.

### Palliative care

Early referral to palliative care can improve quality of life and in some cases survival. Referral should be based on need, not prognosis. For more, visit the Palliative Care Australia website <[www.palliativecare.org.au](http://www.palliativecare.org.au)>.

### Communication

#### The lead clinician and team's responsibilities include:

- discussing treatment options with the patient and/or carer including the intent of treatment as well as risks and benefits
- discussing advance care planning with the patient and/or carer where appropriate
- communicating the treatment plan to the patient's GP
- helping patients to find appropriate support for exercise programs where appropriate to improve treatment outcomes.

### Timeframes

- Surgery should be undertaken **within four weeks** of the MDM, depending on urgency.
- Systemic therapy for unresectable biliary cancer should begin **within four weeks** of the MDM, depending on urgency.
- Adjuvant chemotherapy in postoperative care should begin **within eight weeks** of surgery, depending on the patient's recovery.
- Radiation therapy should begin **within four weeks** of the MDM, depending on urgency.
- Adjuvant radiotherapy in postoperative care should begin **within eight weeks** of surgery, depending on the patient's recovery.

## Step 5: Care after initial treatment and recovery

### Provide a treatment and follow-up summary to the patient, carer and GP outlining:

- the diagnosis, including tests performed and results
- tumour characteristics
- treatment received (types and date)
- current toxicities (severity, management and expected outcomes)
- interventions and treatment plans from other health professionals
- potential long-term and late effects of treatment and care of these
- supportive care services provided
- a follow-up schedule, including tests required and timing

- contact information for key healthcare providers who can offer support for lifestyle modification
- a process for rapid re-entry to medical services for suspected recurrence.

### Communication

#### The lead clinician's responsibilities include:

- explaining the treatment summary and follow-up care plan to the patient and/or carer
- informing the patient and/or carer about secondary prevention and healthy living
- discussing the follow-up care plan with the patient's GP.

### Checklist

- Treatment and follow-up summary provided to the patient and/or carer and the patient's GP.
- Supportive care needs assessment completed and recorded and referrals to allied health services actioned as required.
- Patient-reported outcome measures recorded.

## Step 6: Managing recurrent, residual or metastatic disease

### Detection

Most residual or recurrent disease will be detected via routine follow-up/imaging or by the patient presenting with symptoms.

### Treatment

All patients should be evaluated for referral to the original multidisciplinary team. In managing people with recurrent, residual or metastatic biliary cancer, the treatment aim is to relieve pain and any unnecessary side effects to ensure optimal quality of life.

Treatment will depend on the location and extent of disease, previous management and the patient's preferences.

### Advance care planning

Advance care planning is important for all patients to consider but especially those with advanced disease. It is a voluntary process that allows them to plan for their future health and personal care by thinking about their values and preferences. This can guide future

treatment if the patient is unable to speak for themselves.

### Survivorship and palliative care

Survivorship and palliative care should be addressed and offered early. Early referral to palliative care can improve quality of life and in some cases survival. Referral should be based on need, not prognosis.

### Communication

The lead clinician and team's responsibilities include explaining the treatment intent, likely outcomes and side effects to the patient and/or carer and the patient's GP.

### Checklist

- Treatment intent, likely outcomes and side effects explained to the patient and/or carer and the patient's GP.
- Supportive care needs assessment completed and recorded and referrals to allied health services actioned as required.
- Advance care planning discussed with the patient and/or carer.
- Patient referred to palliative care if appropriate.
- Routine follow-up visits scheduled.

## Step 7: End-of-life care

### Palliative care

Consider a referral to palliative care. Ensure advance care directives have been explained and the patient has been offered an opportunity to develop one if they choose. If the patient has an advance care directive, ensure it is shared with treating teams and is used to guide care decisions if the patient cannot speak for themselves.

### Voluntary assisted dying

Voluntary assisted dying is legislated by state and territory governments, so it is essential to know the rules around this choice in the state or territory where the patient lives.

### Communication

#### The lead clinician's responsibilities include:

- being open about the prognosis and discussing palliative care options with the patient
- establishing transition plans to ensure the patient's needs and goals are considered in the appropriate environment.

### Checklist

- Supportive care needs assessment completed and recorded, and referrals to allied health services actioned as required.
- Patient referred to palliative care.
- Advance care directives (if any) are shared and considered in care decisions.

Visit our guides to best cancer care webpage <[www.cancer.org.au/cancercareguides](http://www.cancer.org.au/cancercareguides)> for consumer guides. Visit our OCP webpage <[www.cancer.org.au/health-professionals/optimal-cancer-care-pathways](http://www.cancer.org.au/health-professionals/optimal-cancer-care-pathways)> for the optimal care pathway and instructions on how to import these guides into your GP software.

# Intent of the optimal care pathways


Optimal care pathways map seven key steps in cancer care. Each of these steps outlines nationally agreed best practice for the best level of care. While the seven steps appear in a linear model, in practice, patient care does not always occur in this way but depends on the particular situation (e.g. the type of cancer, when and how the cancer is diagnosed, prognosis, management, the patient's decisions and their physiological response to treatment).

The principles underpinning optimal care pathways always put patients at the centre of care throughout their experience and prompt the health system to deliver coordinated care.

The optimal care pathways do not constitute medical advice or replace clinical judgement, and they refer to clinical guidelines and other resources where appropriate.

This optimal care pathway can be used in conjunction with population-based optimal care pathways, where appropriate for the person affected by cancer.

Figure 1: The optimal care pathway



## Optimal care pathway resources

There are three resources for each pathway: an optimal care pathway, a quick reference guide for health professionals and a guide to best cancer care for patients, carers and families.


### Optimal care pathways

This optimal care pathway is designed for **health professionals** and **health services**. However, **patients** and **carers** may find useful information in this version to help understand the processes their treating health professionals are following.

This resource aims to:

- assist health professionals to provide optimal care and support to patients with cancer, their families and carers
- provide optimal timeframes for delivering evidence-based care
- emphasise the importance of communication and collaboration between health providers and people affected by cancer
- assist and inform new health professionals or trainees who are entering the cancer care workforce
- provide value to health systems to identify gaps in current cancer services, bring about quality improvement initiatives and improve how services are planned and coordinated. Adherence to the pathways should be measured wherever possible.

Visit the Cancer Council website <<https://www.cancer.org.au/health-professionals/optimal-cancer-care-pathways>> to view the optimal care pathways.



### Quick reference guides

The quick reference guides are for **health professionals** and **health services**. They provide a summary of each optimal care pathway for health professionals and patients.

The quick reference guides include:

- optimal timeframes within which tests or procedures should be completed
- checklists with indicators related to recommendations in the optimal care pathway.

Visit the Cancer Council website <<https://www.cancer.org.au/health-professionals/optimal-cancer-care-pathways>> to view the quick reference guide for this optimal care pathway.



## Guides to best cancer care

The guides to best cancer care are **consumer** resources that help patients understand the optimal cancer care that should be provided at each step. Carers, family and friends may also find the guides helpful.

The guides to best cancer care:

- include optimal timeframes within which tests or procedures should be completed
- include prompt questions to support patients to understand what might happen at each step of their cancer journey and to consider what questions to ask
- provide information to help patients and carers communicate with health professionals
- are available in 11 languages.


Visit the Cancer Council's website <<https://www.cancer.org.au/cancercareguides>> to view the guides to best cancer care.



## Optimal care pathway for Aboriginal and Torres Strait Islander people with cancer

The *Optimal care pathway for Aboriginal and Torres Strait Islander people with cancer* provides a tool to help reduce disparities and improve outcomes and experiences for Aboriginal and Torres Strait Islander people with cancer. This resource can be used in conjunction with the optimal care pathway for each cancer type.


Visit the Cancer Australia website <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/OCP-aboriginal-and-torres-strait-islander-people-cancer-second-edition>> to view the optimal care pathway for Aboriginal and Torres Strait Islander people.



# Principles of the optimal care pathway

The seven principles of care define appropriate and supportive cancer care that is the right of all patients and the right of those caring for and connected with them.

Figure 2: The seven principles underpinning the optimal care pathway



## Principle 1: Patient-centred care



**Patient-centred care informs and involves patients in their care and respects and responds to the preferences, needs and values of patients, families and carers.**

A patient-centred focus increases the experience and satisfaction of patients, their families and carers, and staff, as well as safety and cost-effectiveness (ACSQHC 2019a).

Patient-centred care means:

- patients are informed and involved in decisions about their cancer and the treatment, post-treatment and recovery program ahead
- patients, their families and carers are provided with access to appropriate and accessible health information
- respect for the cultural and religious beliefs of patients and their families is demonstrated when discussing the diagnosis of cancer
- active communication is used to engage patients, their families and carers in the care process – an essential step for patients to be informed
- care processes are mutually beneficial for patients and providers
- special needs are addressed – for example, the needs of people with disabilities or mental health issues.

## **Informed choice and consent**

An informed patient has greater confidence and competence to manage their cancer journey.

Health professionals are responsible for enabling patients to make informed choices according to their preferences, needs and values. Patients should be provided with:

- individualised and timely information and guidance about their treatment
- details of their care, including the advantages and disadvantages of each treatment, the associated potential side effects, the likely outcomes on their performance status (how well a patient is able to carry out activities of daily life) and subsequently their quality of life and any financial implications, at each stage of the pathway (ACSQHC 2020).

Health professionals have a legal responsibility to obtain consent for all procedures from either the patient or their substitute decision-maker if they are not deemed competent.

## **Referral choices and informed financial consent**

Patients have the right to receive the information they need to be able to make an informed decision on where to be referred for treatment. Treating specialists and practitioners should clearly explain the costs or how to find out the costs of services, tests and treatment options upfront to avoid consumers experiencing 'bill shock'.

At the time of referral, the patient's general practitioner or other referring doctor should discuss the different options for referral, waiting periods, expertise, if there are likely to be out-of-pocket costs and the range of services available. This will enable patients to make an informed choice of specialist and health service. Referral decisions influence the care patients receive along the pathway and the direct and indirect costs they and their carers may incur. Different referrals have different costs:

- referral to a public hospital, which may involve some costs
- initial referral to a private specialist with associated costs, with the option of ongoing treatment in a public hospital at any time
- referral to a patient's choice of practitioner for immediate and ongoing private hospital management with associated costs.

Patients should be made aware that even though public hospital health care is 'free' to all Australian citizens and most permanent residents of Australia, there are still associated direct costs such as:

- over-the-counter medication and prescriptions
- wound dressings
- travel costs
- parking fees
- tests that are not covered by Medicare.

A cancer diagnosis and treatment may affect a patient's or carer's income. This is an indirect cost associated with cancer. Social work support is essential to help patients, and their families deal with this issue. Patients should be advised not to undergo private care with significant out-of-pocket expenses if financially constrained. Specialists in private practice need to explain costs at the start of each new treatment to acknowledge the cumulative out-of-pocket expenses that patients can incur.

For more information on informed financial consent refer to Cancer Council's 'Standard for informed financial consent' <<https://www.cancer.org.au/health-professionals/resources/informed-financial-consent>>.

Financial counselling services can provide advice on dealing with financial difficulties. These services can be accessed publicly (via social workers at hospitals, financial counsellors at neighbourhood houses or rural financial aid), privately or through cancer support services such as local charity groups or social work services.

For practical and financial assistance, patients may consider Cancer Council's financial services <<https://www.cancer.org.au/support-and-services/practical-and-financial-assistance>>.

### Shared care

Shared care between a cancer specialist and primary care health professional is delivered in two or more settings by two or more professionals. The primary care provider is usually a general practitioner but can include nurses and allied health practitioners. Shared care can be delivered throughout the care pathway including during treatment, follow-up care, survivorship care and end-of-life care.

Shared care offers several advantages to patients, including the potential for treatment closer to home and more efficient care with less duplication and greater coordination. Evidence comparing shared care and specialised care indicates equivalence in outcomes including recurrence rate, cancer survival and quality of life (Cancer Research in Primary Care 2016).

Telehealth can enable efficient shared care and should be explored for all patients. Patients in some rural or remote locations may access specialists via Medicare Benefit Scheme funded telehealth consultations. General practitioners working in rural or remote locations should be aware of specialist multidisciplinary teams with facilities to reduce the travel burden and costs for patients.

## Principle 2: Safe and quality care



### Hospitals and health professionals are responsible for providing safe and quality care.

Health professionals need to have appropriate training and experience to undertake treatment for biliary cancer. Patients should be referred to an individual practitioner or service with appropriate expertise.

Safe and high-quality care is care provided by appropriately trained and credentialed health professionals who undertake regular quality reviews of their performance, contribute to regular audits of their care and are actively involved in continuing professional development. Hospitals and clinics must have the equipment, staff numbers, policies and procedures in place to support safe and high-quality care for cancer patients. Patients should be offered the safest options for care, which may include using telehealth (Cancer Australia 2020).

Hospital quality committees should ensure all health care is informed by evidence, and health professionals and health service managers (including executives) have a responsibility to evaluate and monitor their practice. Optimal care pathways provide a framework to help evaluate and monitor practice over time. Services should be routinely collecting relevant minimum datasets to support benchmarking, quality care and service improvement. Hospital committees and health professional peak bodies should be auditing this process (ACSQHC 2017, 2020).

The Australian Council on Health Standards <<https://www.achs.org.au/>> has created a set of indicators that helps hospitals conform to appropriate standards.

## Patient-reported experience and outcome measures

Patient-reported experience measures (PREMs) and patient-reported outcome measures (PROMs) should be incorporated into routine cancer care.

PREMs are used to obtain patients' views and observations on aspects of healthcare services they have received (AIHW 2018). Patient experience data is collected for specific services and then relayed to service providers to instigate improvements in patient services (ACSQHC 2019b).

The Australian Hospital Patient Experience Question Set (AHPEQS) is a tool used to assess patient experiences of treatment and care in a private or public hospital. AHPEQS helps to improve the safety and quality of health care by allowing organisations to understand the patient's perspective (ACSQHC 2019b; AIHW 2018).

PROMs measure aspects of a person's health status such as symptoms, quality of life and needs and are collected directly from patients either online, via a smartphone or through paper-based means.

Collecting PROMs, and then instigating an appropriate clinical response, has been shown to prolong survival, reduce health system use and improve patients' quality of life. While there are many sets of PROMs questions that are relevant to any cancer patient, specific questions can be tailored to particular cancer types, populations or different phases of cancer care.

## Principle 3: Multidisciplinary care



**Multidisciplinary care is an integrated team approach that involves all relevant health professionals discussing all relevant treatment options and making joint recommendations about treatment and supportive care plans, taking into account the personal preferences of patients.**

Multidisciplinary care improves patient outcomes. Cancer Australia's 'Principles of multidisciplinary care' provides a flexible definition, allowing services to vary implementation according to cancer type and the service location. The principles stipulate:

- a team approach that involves core disciplines that are integral to providing good care, including general practice, with input from other specialties as required
- communication among team members about treatment planning and plans for follow-up
- access to the full therapeutic range for all patients, regardless of geographical remoteness or size of institution
- care delivery in accordance with nationally agreed standards
- patient involvement in decisions about their care (Cancer Australia 2019).

In addition to these principles, treatment teams should consider clinical trial participation for all eligible patients.

Multidisciplinary meetings, often called MDMs, should be based on the principles outlined above.

For more information on the principles of multidisciplinary care and the benefits of adopting a multidisciplinary approach, refer to Cancer Australia's 'Principles of multidisciplinary care' <<https://www.canceraustralia.gov.au/clinicians-hub/multidisciplinary-care/all-about-multidisciplinary-care/principles-multidisciplinary-care>>.

## Principle 4: Supportive care



**Supportive care is a vital part of any cancer treatment program. Supportive care deals with issues that emerge for patients, families and carers from the effects of the cancer diagnosis and its treatment. It is made up of all the services, information and resources patients may need to meet their physical, psychological, social, information and spiritual needs from the time of diagnosis.**

Supportive care may be 'patient-defined' and based on unmet needs. It is a core component of evidence-based clinical care and its benefits are well established. All cancer patients and their carers should be formally supported and have access to understandable, relevant information about the medical, practical and emotional aspects of the cancer and its treatment (Fitch 2008). The wishes and needs of the patient, their family and their carers should determine the level of support provided. Supportive care is a standard or routine aspect of cancer care and the treatment team should make patients aware of this.

Supportive care should begin from the time of diagnosis and continue throughout the cancer pathway.

For health professionals, supportive care involves:

- screening and assessing patients and families for their supportive care needs
- providing patients with access to a range of multidisciplinary support services, groups and therapies designed to assist them to live with cancer and its treatment and optimise recovery
- optimising referral pathways to community support organisations (cancer-related non-government, not-for-profit and charities) that provide services to cancer survivors – these address many of the care-navigation, psychosocial and information needs of cancer survivors and those affected by cancer (Australian Cancer Survivorship Centre 2021)
- being aware of and delivering culturally appropriate care.
- All members of the multidisciplinary team have a role in providing supportive care along the care pathway, with special attention at transition points.

Supportive care involves routinely and systematically assessing patients to determine their needs. Health professionals can use a variety of validated screening tools for this task (refer to the box below). Clinical review and individual assessment are still required to ensure all patient concerns are identified.



### More information

Visit the WeCan website <[www.wecan.org.au](http://www.wecan.org.au)> for information and resources on supportive care.

### Validated screening tools

- National Comprehensive Cancer Network Distress Thermometer and Problem Checklist
- Supportive Care Needs Assessment Tool for Indigenous People (SCNAT-IP).

## Key review points

The treatment team should assess patients for supportive care needs at these key stages:

- initial presentation or diagnosis (first three months)
- the beginning of treatment or a new phase of treatment
- change in prognosis
- if a patient is found to have a germline genetic mutation predisposing to cancer
- end of treatment
- throughout survivorship
- diagnosis of recurrence
- change in or development of new symptoms
- palliative care
- end-of-life care
- other time points based on clinical judgement.

The team also needs to decide whether the patient requires ongoing referral to supportive care services. Access to services can be through general practice-led chronic disease management plans, team care arrangements and mental health plans. Community support services also have a role to play.

Refer to Appendices A, B and C for more information on supportive care and the specific needs of people that may arise.

## Principle 5: Care coordination



**Care coordination is the responsibility of every professional, both clinical and non-clinical, who works with patients, their families and carers.**

Seamless care coordination is essential for patients to successfully navigate the complex health system. Care coordination is a comprehensive approach to achieving continuity of care for patients. It aims to ensure care is delivered in a systematic, connected and timely way that promotes efficiency and reduces the risk of duplication and over-servicing to meet the medical and personal needs of patients.

Care coordination includes:

- proactive and timely communication with patients, their families and carers
- treatment plans, survivorship care plans and/or advance care planning
- coordinated appointments to ensure timely diagnosis, treatment and survivorship care
- appropriate tests and results being available to the treating team so treatment decisions can be made
- medical records being available to all members of the treating team and at scheduled appointments
- translation or interpreter services arranged if the patient/carer is from a non-English-speaking background or has difficulty communicating due to a physical disability
- practical support such as transport, accommodation, advance care planning and financial support
- referral and access to supportive care
- access to clinical trials
- access to telehealth for people in rural and remote areas and for managing vulnerable patients.

Care coordination brings together different health professionals, teams and health services. It also encompasses MDMs, multidisciplinary assessment clinics, supportive care screening and assessment, referral practices, data collection, common protocols, information for patients and individual clinical treatment.

Care coordination should cross the acute and primary care interface and should aim to achieve consistency of care through clear communication, linkages and collaborative integrated care planning.

Care coordination can be facilitated through electronic health record management such as My Health Record. My Health Record is a secure online database that helps with data collection and care coordination (My Health Record 2019).

Formal care coordination through appointed care coordinators plays an important role in managing and supporting patients through the health system. The availability of dedicated care coordinators varies across states and territories according to the complexity of care required and local service capacity and resourcing.

## Principle 6: Communication



**Everyone employed in the health system is responsible for ensuring the communication needs of patients, their families and carers are met.**

Good and open communication is a key principle of care for cancer patients. This includes communication between oncology and primary care health professionals and with patients. General practitioners should be involved in care from the point of diagnosis, and patients should be encouraged to maintain a relationship with their general practitioner through all stages of cancer care. Communication should be regular and timely.

Attendance of a family member or carer at clinical appointments is beneficial for many patients, as the family member or carer can provide informational and emotional support. General practitioners and clinicians should encourage and support the involvement of family members and carers by providing an inclusive and supportive consultation environment (Laidsaar-Powell et al. 2018a). Laidsaar-Powell et al. provide evidence-based guidance on how to support family member or carer involvement in consultations (Laidsaar-Powell et al. 2018a, 2018b).

Every person with cancer will have different communication needs, including cultural and language differences. When anyone involved in treatment communicates with patients, they should be truthful and transparent but aware of cultural and psychological sensitivities. In communicating with patients, healthcare providers should undertake to:

- empower patients to be active in treatment discussions
- use professionally trained interpreters if required – for example, when communicating with people from culturally diverse backgrounds whose primary spoken language is not English and for people with a hearing impairment (visit the Translating and Interpreting Services website <[www.tisnational.gov.au](http://www.tisnational.gov.au)> for more information on interpreter and language services)
- use culturally sensitive and appropriate forums of communication for people from culturally diverse backgrounds and Aboriginal and Torres Strait Islander people, as appropriate
- provide appropriate information for people from culturally diverse backgrounds
- provide information on community-based supportive care services and resources to patients and their families and carer

- identify the patient's substitute treatment decision-maker to ensure they are involved in relevant discussions
- ensure patients, their families or their carers have the opportunity to ask questions
- seek consent before conveying information between health professionals or healthcare teams or with family and carers
- be respectful if a patient seeks a second opinion from another health professional
- ensure patients do not have to convey information between areas of care (it is the provider's and health system's responsibility to transfer information between areas of care)
- communicate in plain language (avoiding complex medical terms and jargon)
- ensure information is communicated at a level relevant to the patient's health literacy and that of their families and carers (ACSQHC 2020)
- use tools, diagrams and aids as appropriate (Gilligan et al. 2017)
- ensure the patient is aware of how to access electronic patient information, where appropriate
- allow enough time for communication, especially when conveying complex or sensitive information such as an initial diagnosis
- check the patient's and/or their family or carer's understanding by asking the patient and/or their family or carer to say in their own words what has been conveyed.

Healthcare providers should also consider offering patients a question prompt list before a consultation and recordings or written summaries of their consultations afterwards. Question prompt lists are effective in improving communication and the psychological and cognitive outcomes of cancer patients. Recordings or summaries of key consultations improve patients' recall of information and satisfaction (Hack et al. 2012). Written care plans, treatment summaries, survivorship care plans and advance care directives are effective records and communication tools.

Communication skills training programs that use role-play to develop skills and observe patient interactions to provide feedback, should be available to health professionals at every level of practice (Gilligan et al. 2017).

Communication skills training programs and resources can be found on the following websites:

- Australian Commission on Safety and Quality in Healthcare, Communicating for safety resource portal <<https://c4sportal.safetyandquality.gov.au>>
- state and territory Cancer Councils <[www.cancer.org.au/about-us/state-and-territory-councils/](http://www.cancer.org.au/about-us/state-and-territory-councils/)> for the relevant council
- eviQ <<https://education.eviq.org.au>>
- VITAL talk <[www.vitaltalk.org](http://www.vitaltalk.org)>
- Pam McLean Centre <<https://pammcleancentre.org>>.

Telehealth has become an increasingly acceptable alternative to face-to-face consultations. When using telehealth, the team must consider what is best for the patient, including the patient's preferences of the patient. A face-to-face consultation should be the first option, if it is safe, when delivering critical diagnosis information, a change in therapy or prescribing intensive treatment. If this is not an option, a video consultation should be considered, and the patient should be encouraged to have a support person with them to assist (Cancer Australia 2020).

## Principle 7: Research and clinical trials



**Research and clinical trials play an important role in establishing the efficacy and safety of diagnostic, prognostic and therapeutic interventions, as well as establishing the role of psychological, supportive care and palliative care interventions (Sjoquist & Zalcborg 2013).**

Clinical trials are the foundation for improved cancer outcomes, allowing new treatments to be tested and offering patients access to potentially more effective therapies than otherwise available to them.

Clinical trials are available for multiple types of cancer and may be a valuable option for people with rare, difficult-to-treat conditions for which there may be limited evidence about how the condition is best treated or managed (Australian Clinical Trials 2015).

Treating specialists and multidisciplinary teams should be aware of or search for clinical trials that may be suitable for their patients. Specialists should be willing to refer appropriate patients to other treating centres to participate in research or clinical trials at any stage of the care pathway and be willing to discuss the pros and cons of participating in such trials. Any member of the multidisciplinary team can encourage cross-referral between clinical trials centres. Possible ineligibility to participate in a clinical trial should be discussed with the patient. Acknowledge disappointment and offer support in this instance.

Health services should strive to implement policies and procedures that facilitate equitable access to clinical trials for all patients, including culturally diverse patients, regional patients and those from Aboriginal or Torres Strait Islander communities.

The use of telehealth technology, such as the Australasian Tele-trial Model, hopes to improve access to trials for patients being treated in rural and regional areas (COSA 2016).

Australian Cancer Trials is a national clinical trials database. It provides information on the latest clinical trials in cancer care, including trials that are recruiting new participants. Search for a trial <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)> via its website.

### **Education and training**

Research and clinical trials provide an opportunity to educate health professionals who are in training. Cancer centres may be affiliated with teaching hospitals, universities or research groups to promote higher education or to develop the academic workforce, leading to more sustainable practice. Specialists should be encouraged to take up and retain active membership to professional societies and organisations that can assist with professional development opportunities.

## Summary – optimal timeframes

Evidence-based guidelines, where they exist, should inform timeframes. Treatment teams need to recognise that shorter timeframes for appropriate consultations and treatment can promote a better experience for patients. Three steps in the pathway specify timeframes for care (Figure 3). They are designed to help patients understand the timeframes in which they can expect to be assessed and treated, and to help health services plan care delivery in accordance with expert-informed time parameters to meet the expectation of patients. These timeframes are based on expert advice from the Biliary Cancer Expert Working Group. The pathway from presentation with symptoms at a general practice to treatment should take no longer than 10 weeks. Different stages of the pathway may overlap according to local practice.

Figure 3: Timeframes for care

Step in pathway	Care point	Timeframe
Presentation, initial investigations and referral	Signs and symptoms	Presenting symptoms should be <b>immediately</b> assessed by a GP and investigations begun.
	Initial investigations initiated by GP	Patients who present with jaundice should be referred for tests <b>immediately</b> and followed up rapidly. Other symptoms require investigations to be completed and reviewed by GP <b>within 2 weeks</b> .
	Referral to specialist	Patients with suspected or proven biliary cancer should have immediate referral and be seen by a specialist linked to a multidisciplinary team <b>within 2 weeks</b> of referral.
Diagnosis, staging and treatment planning	Diagnosis and staging	Diagnostic and staging investigations should be completed <b>within 2 weeks</b> of referral and can occur concurrently with other steps in the pathway.
	Multidisciplinary team meeting and treatment planning	The patient case must be referred to an MDM <b>within 2 weeks</b> of finding a suspected biliary cancer. The MDM can occur after the minimum diagnostic tests. The full range of tests should not delay presentation at an MDM.
Treatment	Surgery	Surgery should be undertaken <b>within 4 weeks</b> of MDM, depending on urgency and modality.
	Systemic therapy	Systemic therapy for unresectable biliary cancer should begin <b>within 4 weeks</b> of the MDM, depending on urgency and modality. Adjuvant chemotherapy in postoperative care should begin <b>within 8 weeks</b> of surgery, depending on the patient's recovery.
	Radiation therapy	Radiation therapy should begin <b>within 4 weeks</b> of MDM, depending on urgency and modality. Adjuvant radiotherapy in postoperative care should begin <b>within 8 weeks</b> of surgery, depending on the patient's recovery.

# Optimal care pathway

## Seven steps of the optimal care pathway

Step 1: Prevention and early detection

Step 2: Presentation, initial investigations and referral

Step 3: Diagnosis, staging and treatment planning

Step 4: Treatment

Step 5: Care after initial treatment and recovery

Step 6: Managing recurrent, residual or metastatic disease

Step 7: End-of-life care

This optimal care pathway outlines a guide to best practice for biliary cancer. Biliary cancer includes gallbladder cancer and biliary tract cancer. Biliary tracts are a system of ducts that drain bile and other substances from the liver to the small bowel. Biliary tract cancer is also called cholangiocarcinoma. Biliary cancer is rare. In Australia, around 2,226 people were diagnosed with either gallbladder or biliary tract cancer in 2023, although incidence is increasing (GI Cancer Institute 2024). On average, five-year survival for biliary cancer across all ages is 20 per cent (GI Cancer Institute 2024). The average age at diagnosis is 72 years.

Most biliary cancers are adenocarcinomas – tumours that arise from the gland cells of the tissues lining the bile ducts or the gallbladder. Biliary tract cancer can form anywhere along the bile ducts, and each sub-type of biliary tract cancer is named after the location where the primary cancer begins. Intrahepatic biliary tract cancer forms in the bile ducts inside the liver. Intrahepatic biliary tract cancers are also called intrahepatic cholangiocarcinomas. Extrahepatic biliary tract cancer forms in the bile ducts outside the liver. The two major types of extrahepatic biliary tract cancer are perihilar biliary tract cancer and distal biliary tract cancer:

- **Perihilar biliary tract cancer** is found in the area where the right and left bile ducts (ducts draining the right lobes and the left lobes of the liver respectively) exit the liver and join to form the common hepatic duct. Perihilar biliary tract cancer is also called a Klatskin tumour or perihilar cholangiocarcinoma.
- **Distal biliary tract cancer** is found in the area of the common bile duct as it passes through the pancreas and ends in the small intestine. Distal biliary tract cancer is also called extrahepatic cholangiocarcinoma.

Determining the incidence of the subtypes of biliary cancer is problematic because there is no specific ICD code for each subtype, a coding system that ensures accuracy in understanding trends in all aspects of the disease (Khan et al. 2019). Historically these cancers have not been well-recognised and have often been labelled as liver cancer, cancer of unknown primary and pancreatic cancer, further contributing to inaccurate data (Khan et al. 2019).

Despite the slight increase in survival, biliary cancer is a low-survival cancer. Even if there are good initial treatment outcomes, the recurrence rate is high. Given the poor prognosis of this cancer, even when treatment is given with curative intent, poor outcomes may result. Early specialist palliative care will be required for patients with biliary cancer, especially at the time of recurrence or spread to other organs.

More information about managing periampullary cancer and distal cholangiocarcinoma found in the pancreas is in the pancreatic cancer optimal care pathway.

## Step 1: Prevention and early detection

This step outlines recommendations for the prevention and early detection of biliary cancer.

Evidence shows that not smoking, avoiding or limiting alcohol intake, eating a healthy diet, maintaining a healthy body weight, being physically active, being sun smart and avoiding exposure to oncoviruses or carcinogens may help reduce cancer risk (Cancer Council Australia 2024).

### 1.1 Prevention

The causes of biliary cancer are not fully understood, and there is currently no clear prevention strategy.

### 1.2 Risk factors

Gallbladder cancer	Biliary tract cancer
<ul style="list-style-type: none"><li>• Having gallstones</li><li>• Having gallbladder polyps</li><li>• Medical history of long-term inflammation of the gallbladder</li><li>• History of a first-degree or immediate relative (mother, father, sibling or child) having gallbladder cancer can increase the risk five-fold, but given that gallbladder cancer is extremely rare, the risk is small</li><li>• Inherited genetic conditions such as abnormalities of the liver, gallbladder and bile ducts</li><li>• Being overweight or obese</li><li>• Smoking</li><li>• Being from East Asia or South America (Baria et al. 2022)</li></ul>	<ul style="list-style-type: none"><li>• Medical history of long-term inflammation of the bile duct such as primary sclerosing cholangitis</li><li>• Biliary cysts</li><li>• Inherited genetic conditions such as abnormalities of the liver, gallbladder and bile ducts</li><li>• Chronic liver disease such as cirrhosis, hepatitis B virus infection and hepatitis C virus infection</li><li>• Infection with a liver fluke parasite has been linked to bile duct cancer cases in Africa and Asia</li><li>• Genetic changes such as hereditary nonpolyposis colorectal cancer syndrome – Lynch syndrome, characterised by abnormalities of the mismatch repair genes (MMR) <i>MLH1</i>, <i>MSH2</i>, <i>MSH6</i> and <i>PMS2</i>, as well as mutations in <i>BRCA1/2</i>.</li><li>• Exposure to Thorotrast, which is a radioactive compound that was used in x-rays from the 1930s to the 1950s</li><li>• Exposure to asbestos</li><li>• Being from East Asia or South America (Baria et al. 2022)</li><li>• Pre-existing inflammatory bowel disease such as chronic ulcerative colitis</li><li>• Older age</li><li>• Diabetes</li><li>• Obesity</li></ul>

### 1.3 Early detection

#### 1.3.1 Screening recommendations

Routine screening for biliary cancer is not currently recommended in either the general population or in relatives of people with biliary cancer.

For people at higher risk, such as those with primary sclerosing cholangitis, long-term inflammation of the gallbladder or gallbladder polyps (6–10 mm), consider annual surveillance using ultrasound. Consider cholecystectomy for patients with gallbladder polyps over 10 mm.

## Step 2: Presentation, initial investigations and referral

**This step outlines the process for the general practitioner to initiate the right investigations and refer to the appropriate specialist in a timely manner. The types of investigations the general practitioner undertakes will depend on many factors, including access to diagnostic tests, the availability of medical specialists and patient preferences.**

### 2.1 Signs and symptoms

Many cases present with non-specific symptoms or are asymptomatic until at advanced stages of the disease. The following signs and symptoms should be investigated:

- jaundice with or without darkened urine, pale bowel movements and itchy skin
- unexplained nausea and vomiting
- unexplained weakness
- unexplained loss of appetite and weight loss
- fevers and chills
- pain in the right side of the abdomen.

The presence of multiple signs and symptoms, particularly in combination with other underlying risk factors, indicates an increased risk of biliary cancer.

#### 2.1.1 Timeframe for general practitioner consultation

Presenting symptoms should be immediately assessed by a general practitioner and investigations begun.

### 2.2 Assessments by the general practitioner

Depending on the presenting symptoms and risk factors, the general practitioner investigations include the following.

Where there is suspicion of biliary cancer, consider:

- an abdominal CT scan and/or abdominal ultrasound
- serum CA 19-9, liver function tests, full blood examination, biochemistry and coagulation screen (INR/APTT)
- early referral to a specialist doctor (strongly indicated) (gastroenterologist, hepatologist, oncologist or hepatopancreatobiliary [HPB] surgeon) with expertise in biliary cancer management and access to a multidisciplinary team, usually before a definitive diagnosis is made.

Where jaundice is present, the following should be performed, reviewed and the patient referred to a specialist doctor urgently:

- liver function tests and coagulation screen (INR/APTT)
- abdominal ultrasound
- CT of the abdomen and pelvis including a multiphase scan of the liver, provided this does not delay urgent referral.

An abnormal result should be discussed face to face with the patient and information provided.

### 2.2.1 Timeframe for completing investigations

Patients who present with jaundice should be referred for tests immediately and followed up rapidly. Other symptoms require investigations to be completed and reviewed by a general practitioner **within two weeks**.

## 2.3 Initial referral

If the cancer diagnosis is confirmed, suspected, or the results are inconsistent or indeterminate, the general practitioner must refer the patient to an appropriate specialist (gastroenterologist, hepatologist, oncologist or HPB surgeon) with expertise in biliary cancer management and access to a multidisciplinary team, to complete the diagnosis. In patients with locally advanced, non-metastatic disease, if resectability needs to be determined, referral should be to an HPB surgeon who works as part of a multidisciplinary team.

A cancer nurse coordinator (upper gastrointestinal cancer nurse coordinator if available) should be involved in the care and coordination of the patient from referral to specialist and MDM.

Patients should be enabled to make informed decisions about their choice of specialist and health service. General practitioners should make referrals in consultation with the patient after considering the clinical care needed, cost implications (refer to referral options and informed financial consent on page 7), waiting periods, location and facilities, including discussing the patient's preference for health care through the public or the private system.

Referral for suspected or diagnosed biliary cancer should include the following essential information to accurately triage and categorise the level of clinical urgency:

- important psychosocial history and relevant medical history
- family history, current symptoms, medications and allergies
- results of current clinical investigations (imaging and pathology reports)
- results of all prior relevant investigations
- notification if an interpreter service is required.

Many services will reject incomplete referrals, so it is important that referrals comply with all relevant health service criteria.

If access is via online referral, a lack of a hard copy should not delay referral.

The specialist should provide timely communication to the general practitioner about the consultation and should notify the general practitioner if the patient does not attend appointments.

Aboriginal and Torres Strait Islander patients will need a culturally appropriate referral. To view the optimal care pathway for Aboriginal and Torres Strait Islander people with cancer and the corresponding quick reference guide, visit the Cancer Australia website <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/OCP-aboriginal-and-torres-strait-islander-people-cancer-second-edition>>. Download the consumer resources – *Checking for cancer* and *Cancer* from the Cancer Australia website <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/checking-cancer-what-expect>> and the Cancer Australia website <<https://www.canceraustralia.gov.au/impacted-cancer>>.

### 2.3.1 Timeframe for referring to a specialist

Where there is a confirmed diagnosis or a high level of suspicion, patients should be seen by a specialist (gastroenterologist, hepatologist, oncologist or HPB surgeon) with expertise in biliary cancer management and linked to a multidisciplinary team **within two weeks** of referral to the specialist.

The multidisciplinary team should have a rapid access program or contact person to set up the appointment. This is commonly the nurse coordinator for the service.

## 2.4 Support and communication

### 2.4.1 Supportive care

The patient's general practitioner should consider an individualised supportive care assessment where appropriate to identify the needs of an individual, their carer and family. Refer to appropriate support services as required. Refer to the validated screening tools mentioned in Principle 4 'Supportive care'.

A number of specific needs may arise for patients at this time:

- assistance for dealing with the emotional distress and/or anger of dealing with a potential cancer diagnosis, anxiety/depression, interpersonal problems and adjustment difficulties
- management of physical symptoms including pain, fatigue, nausea, vomiting, poor appetite, weight loss and itch
- nutritional support if weight loss is significant – refer to an accredited dietitian for assessment and support
- encouragement and support to increase levels of exercise (Cormie et al. 2018; Hayes et al. 2019) – consider referral to an accredited practising exercise physiologist or physiotherapist.

For more information refer to the National Institute for Health and Care Excellence 2015 guidelines, *Suspected cancer: recognition and referral* <[www.nice.org.uk/guidance/ng12/chapter/Recommendations-on-patient-support-safety-netting-and-the-diagnostic-process](http://www.nice.org.uk/guidance/ng12/chapter/Recommendations-on-patient-support-safety-netting-and-the-diagnostic-process)>.

For additional information on supportive care and needs that may arise for different population groups, refer to Appendices A, B and C.

### 2.4.2 Communication with patients, carers and families

The general practitioner is responsible for:

- providing patients with information that clearly describes to whom they are being referred, the reason for referral and the expected timeframes for appointments
- requesting that patients notify them if the specialist has not been in contact within the expected timeframe
- considering referral options for patients living rurally or remotely
- supporting the patient while waiting for the specialist appointment (Cancer Council nurses are available to act as a point of information and reassurance during the anxious period of awaiting further diagnostic information; patients can contact 13 11 20 nationally to speak to a cancer nurse).



#### More information

Refer to Principle 6 'Communication' for communication skills training programs and resources.

## Step 3: Diagnosis, staging and treatment planning

**Step 3 outlines the process for confirming the diagnosis and stage of cancer and for planning subsequent treatment. The guiding principle is that interaction between appropriate multidisciplinary team members should determine the treatment plan.**

### 3.1 Specialist diagnostic work-up

The treatment team, after taking a thorough medical history and making a thorough medical examination of the patient, should undertake the following investigations under the guidance of a specialist.

The minimum diagnostic procedure that should be completed before the MDM is a CT scan. The other tests can be performed under the direction of the MDM and should not delay discussing a patient with suspected biliary cancer to the MDM. If the remaining staging tests reveal contrary information the patient may be re-presented to the MDM for further consideration.

Diagnosis of a mass is primarily by imaging. Contrary to the situation with most cancers, definitive tissue diagnosis before multidisciplinary planning is not recommended. Biopsy approaches may be complex but are generally required where there is diagnostic uncertainty or where tissue is required for systemic therapy (e.g. chemotherapy) or clinical trials. This should be facilitated and overseen by the MDM process. Final pathology may, rarely, prove benign, despite appropriate work-up.

Contrast-enhanced CT scan of the chest, abdomen and pelvis, including a multiphase CT scan of the liver, is recommended (to be completed first if not already performed).

The following investigations are suggested and can occur under the direction of the MDM:

- contrast-enhanced MRI liver and MRCP
- endoscopic ultrasound (EUS)
- FDG-PET. (This can be useful in some cases but is not necessary in all cases. It may be a useful tool, for example, to evaluate evidence of spread of intrahepatic tumours or equivocal findings on radiology.)

The treatment team may consider asking an interventional endoscopist to perform endoscopic retrograde cholangiopancreatography (ERCP) with or without SpyGlass cholangioscopy to further confirm the diagnosis or in equivocal cases.

#### **Special consideration: stenting**

In cases where there is biliary obstruction at the initial presentation stage, carefully consider the role and timing of stenting. Urgent stenting is not recommended if the patient's symptoms are mild. Appropriate diagnostic work-up should take place first, and wherever possible the initial insertion of a stent should be managed through the MDM.

At a minimum, a CT scan with appropriately selected arterial and venous contrast protocols with or without an MRI liver/MRCP should be performed to clearly identify the resectability status **before** stenting and other interventions. The information obtained from imaging will determine the type of stent (e.g. plastic versus metal) and how stenting should be performed (e.g. via ERCP, sometimes combined with EUS, or during percutaneous transhepatic cholangiography [PTC]).

Stenting should not be done before the patient is reviewed at an MDM, and the drainage approach taken is highly dependent on local expertise.

The choice of stent is made on a case-by-case basis. The approximate life of a plastic stent, which is more likely to block, is three months, while a metal stent lasts up to a year. The choice of stent should therefore be based on how long the patient is expected to need the stent before further intervention and what other interventions are planned.

### 3.1.1 Timeframe for completing investigations

Diagnostic investigations should be completed **within two weeks** of referral and can occur alongside other steps in the pathway.

### 3.1.2 Genetic testing (family risk)

Inherited genetic conditions causing biliary cancer are uncommon. The types of genetic changes that are associated with an increased risk of biliary cancer include hereditary nonpolyposis colorectal cancer syndrome (Lynch syndrome), characterised by abnormalities of the mismatch repair genes (MMR) *MLH1*, *MSH2*, *MSH6*, and *PMS2*, as well as mutations in *BRCA1/2*.

Biliary tumours should be routinely checked for mutations in the MMR genes and, if present, test the patient's germline DNA for inherited abnormalities.

Patients with a strong family history of cancer should be referred to a family cancer clinic to determine whether the tumour is part of an inherited predisposition to cancer.

In some cases certain pathological subtypes of cancer or tumour tests (immunohistochemistry or tumour genetic tests) may suggest an underlying inherited cancer predisposition.

Anyone diagnosed with cancer should have a detailed personal and family cancer history taken. Consult relevant guidelines <<https://www.eviq.org.au/cancer-genetics/referral-guidelines>> to determine if referral to a familial cancer service is appropriate.

A familial cancer service assessment can determine if genetic testing is appropriate. Genetic testing is likely to be offered when there is at least a 10 per cent chance of finding a causative 'gene error' (pathogenic gene variant; previously called a mutation). Usually testing begins with a variant search in a person who has had cancer (a diagnostic genetic test). If a pathogenic gene variant is identified, variant-specific testing is available to relatives to see if they have or have not inherited the familial gene variant (predictive genetic testing).

These sorts of genetic tests should be distinguished from molecular analysis of the biliary cancer to determine if there are any acquired genetic abnormalities that may lend themselves to targeted drug therapy.

Medicare funds some genetic tests via a Medicare Benefits Schedule (MBS) item number but most are not. Depending on the personal and family history, the relevant state health system may fund public sector genetic testing.

Pre-test counselling and informed consent is required before any genetic testing. In some states the treating team can offer 'mainstream' diagnostic genetic testing, after which referral is made to a familial cancer service if a pathogenic gene variant is identified. The familial cancer service can provide risk management advice, facilitate family risk notification and arrange predictive genetic testing for the family.

Visit the Centre for Genetics Education website <<https://www.genetics.edu.au/SitePages/Genomic-Testing-Family-Considerations.aspx>> for basic information about cancer in a family.

### 3.1.3 Pharmacogenetics

Pharmacogenetics describes how individual genetic differences can lead to differences in the way certain medicines interact with the body. These interactions can affect the effectiveness of medications and any side effects. Applying pharmacogenetics to treatment planning may help patients to be prescribed the most appropriate treatment at the optimal dose from the beginning of treatment (NHMRC 2013).

## 3.2 Staging

Staging is a critical element in treatment planning and should be clearly documented in the patient's medical record. Staging for biliary cancer is vital to determine if the disease is resectable or non-resectable because the treatment pathways are different. Biliary cancer is considered resectable if it can be entirely removed by surgery. Staging and discussion at an MDM must have been completed to undertake a curative resection. Staging parameters include radiological imaging (tumour size, number and location of lesions, presence or otherwise of metastases and vascular invasion) and Eastern Cooperative Oncology Group (ECOG) status, which should be documented to help determine whether patients are fit for various procedures (e.g. surgery or chemotherapy).

### 3.2.1 Resectable biliary cancer

Biopsy may not be necessary if the multidisciplinary team is planning a curative resection and there is a high suspicion of biliary cancer. Staging is carried out according to the American Joint Committee on Cancer's TNM staging system to guide the treatment pathway and is specific to every subtype of biliary cancer.

Perihilar biliary cancers are further classified according to the Bismuth-Corlette classification to describe their anatomical location. This guides the surgical pathway.

The staging evaluation of patients with suspected resectable biliary cancer is best accomplished using a selection of the following studies:

- contrast-enhanced CT of the chest, abdomen and pelvis, including multiphase CT liver
- contrast-enhanced MRI liver and MRCP
- consideration of PET scan
- consideration of EUS
- serum CA 19-9 – while not diagnostic, it can be useful for prognosis, liver function tests, full blood examination, biochemistry and coagulation screen.

Diagnostic laparoscopy may provide additional staging information for tumours that are considered resectable (to exclude unsuspected peritoneal metastases). Percutaneous or laparoscopic biopsy of the primary tumour is not recommended in patients who may be candidates for liver transplantation because there is a risk of disseminating disease following those procedures. Endoscopic biopsy with EUS, ERCP or SpyGlass cholangiography may be a reasonable strategy. A presumed radiographic diagnosis of biliary cancer is sufficient in noncirrhotic patients in whom a decision has been made to proceed with surgical resection.

### 3.2.2 Non-resectable biliary cancer

For every patient where there are concerns of local extension of the tumour, with suspected vascular involvement and/or hepatic metastasis, the ideal investigations are:

- contrast-enhanced CT scan of the chest, abdomen and pelvis, including multiphase CT of the liver
- contrast-enhanced MRI liver and MRCP
- FDG-PET (may help identify nodal and distant metastases).

#### Special consideration: biopsy

In the case of unresectable disease (confirmed at an MDM), a core biopsy should be obtained for diagnostic pathology and molecular profiling before systemic or locoregional therapies are started. Pathological diagnosis is based on the WHO classification for biliary cancer. Molecular characterisation of biliary cancer may be used to guide treatment in advanced disease.

In undertaking biopsy, an individualised approach that considers tumour type, resectability, location, extent, presence or absence of jaundice, patient performance status (ECOG status) and local expertise is optimal and should consider the following:

- Core biopsy is preferable to brush cytology.
- Core biopsy is preferable to fine needle aspiration.
- Depending on location, EUS-guided core biopsy may be an option to obtain biopsies of enlarged regional nodes and to obtain a tumour biopsy if ERCP-guided biopsies are negative or inconclusive or a stent is not required.
- Gastroenterologists should be encouraged to attempt ERCP with biliary forceps, once this approach has been recommended by the MDM.
- ERCP with biliary forceps is not recommended in undrained parts of the liver.
- Percutaneous transabdominal, and transduodenal, approaches to the hilum should not be used for resectable or transplantable disease due to the risk of seeding and sepsis.
- Biopsy of hilar lesions via PTC or endoscopically (from within the biliary tree) at ERCP may be appropriate.
- For most cases, review by a high-volume centre is recommended.

### 3.2.3 Timeframe for completing investigations

Staging investigations that have been recommended by the MDM should be completed **within two weeks** of referral.



#### More information

Visit the Cancer Institute New South Wales website for information about understanding the stages of cancer <[www.cancer.nsw.gov.au/understanding-cancer/stages-of-cancer](http://www.cancer.nsw.gov.au/understanding-cancer/stages-of-cancer)>.

## 3.3 Performance status

Patient performance status is a central factor in cancer care and should be clearly documented in the patient's medical record.

Performance status should be measured and recorded using an established scale such as the ECOG scale.

## 3.4 Treatment planning

### 3.4.1 Key considerations beyond treatment recommendations

A number of factors should be considered at this stage:

- the patient's overall condition, life expectancy, personal preferences and decision-making capacity
- discussing the multidisciplinary team approach to care with the patient
- appropriate and timely referral to an MDM
- pregnancy and fertility
- support with travel and accommodation
- teleconferencing or videoconferencing as required.

### 3.4.2 Timing for multidisciplinary team planning

The multidisciplinary team should meet to discuss newly diagnosed patients before definitive treatment so that a treatment plan can be recommended and there can be early preparation for the post-treatment phase. Patients with proven or suspected biliary cancer should be discussed at an MDM **within two weeks** of the initial specialist review. The MDM can occur after the minimum diagnostic tests, the full range of tests should not delay presentation at an MDM.

The level of discussion may vary, depending on the patient's clinical and supportive care factors. Very occasionally, some patients with non-complex cancers may not be discussed by a multidisciplinary team; instead the team may have treatment plan protocols that will be applied if the patient's case (cancer) meets the criteria. If patients are not discussed at an MDM, they should at least be named on the agenda for noting. The proposed treatment must be recorded in the patient's medical record and should be recorded in an MDM database where one exists.

Teams may agree on standard treatment protocols for non-complex care, facilitating patient review (by exception) and associated data capture.

Results of all relevant tests and access to images should be available for the MDM. Information about the patient's concerns, preferences and social and cultural circumstances should also be available.

### 3.4.3 Responsibilities of the multidisciplinary team

The multidisciplinary team requires administrative support in developing the agenda for the meeting, for collating patient information and to ensure appropriate expertise around the table to create an effective treatment plan for the patient. The MDM has a chair and multiple lead clinicians. Each patient case will be presented by a lead clinician (usually someone who has seen the patient before the MDM). In public hospital settings, the registrar or clinical fellow may take this role. A member of the team records the outcomes of the discussion and treatment plan in the patient history and ensures these details are communicated to the patient's general practitioner. The team should consider the patient's values, beliefs and cultural needs as appropriate to ensure the treatment plan is in line with these.

#### **3.4.4 Members of the multidisciplinary team for biliary cancer**

The multidisciplinary team should be composed of the core disciplines that are integral to providing good care. Team membership should reflect both clinical and supportive care aspects of care. Pathology and radiology expertise are essential.

Refer to Appendix E for a list of team members who may be included in the multidisciplinary team for biliary cancer.

Core members of the multidisciplinary team are expected to attend most MDMs either in person or remotely via virtual mechanisms. Additional expertise or specialist services may be required for some patients. An Aboriginal and Torres Strait Islander cultural expert should be considered for all patients who identify as Aboriginal or Torres Strait Islander.

#### **3.4.5 Responsibilities of individual team members**

The general practitioner who made the referral is responsible for the patient until care is passed to another practitioner who is directly involved in planning the patient's care.

The general practitioner may play a number of roles in all stages of the cancer pathway including diagnosis, referral, treatment, shared follow-up care, post-treatment surveillance, coordination and continuity of care, as well as managing existing health issues and providing information and support to the patient, their family and carer.

A nominated contact person from the multidisciplinary team may be assigned responsibility for coordinating care in this phase. Care coordinators are responsible for ensuring there is continuity throughout the care process and coordination of all necessary care for a particular phase (COSA 2015). The care coordinator may change over the course of the pathway.

The lead clinician is responsible for overseeing the activity of the team and for implementing treatment within the multidisciplinary setting.

### **3.5 Research and clinical trials**

Patients should be encouraged to participate in research or clinical trials where available and appropriate. Clinical trials accelerate knowledge gathering and can provide options for cancer patients when other therapies have failed or are yet to be approved. Genomics clinical trials may offer matched therapies not yet available as part of routine care.

For more information visit the Cancer Australia website <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>.

## **3.6 Support and communication**

### **3.6.1 Prehabilitation**

Cancer prehabilitation uses a multidisciplinary approach combining exercise, nutrition and psychological strategies to prepare patients for the challenges of cancer treatment such as surgery, systemic therapy and radiation therapy. Team members may include anaesthetists, oncologists, surgeons, haematologists, clinical psychologists, exercise physiologists, physiotherapists and dietitians, among others.

Patient performance status is a central factor in cancer care and should be frequently assessed. All patients should be screened for malnutrition using a validated tool, such as the Malnutrition Screening Tool (MST). The lead clinician may refer obese or malnourished patients to a dietitian preoperatively or before other treatments begin.

Patients who currently smoke should be encouraged to stop smoking before receiving treatment. This should include an offer of referral to Quitline in addition to smoking cessation pharmacotherapy if clinically appropriate.

Evidence indicates that patients who respond well to prehabilitation may have fewer complications after treatment. For example, those who were exercising before diagnosis and patients who use prehabilitation before starting treatment may improve their physical or psychological outcomes, or both, and this helps patients to function at a higher level throughout their cancer treatment (Cormie et al. 2017; Silver 2015).

For patients with biliary cancer, the multidisciplinary team should consider these specific prehabilitation assessments and interventions for treatment-related complications or major side effects:

- conducting a physical and psychological assessment to establish a baseline function level
- identifying impairments and providing targeted interventions to improve the patient's function level (Silver & Baima 2013)
- reviewing the patient's medication to ensure optimisation and to improve adherence to medicine used for comorbid conditions.

Following completion of primary cancer treatment, rehabilitation programs have considerable potential to enhance physical function.

### 3.6.2 Fertility preservation and contraception

Cancer and cancer treatment may cause fertility problems. This will depend on the age of the patient, the type of cancer and the treatment received. Infertility can range from difficulty having a child to the inability to have a child. Infertility after treatment may be temporary, lasting months to years, or permanent (AYA Cancer Fertility Preservation Guidance Working Group 2014).

Patients need to be advised about and potentially referred for discussion about fertility preservation before starting treatment and need advice about contraception before, during and after treatment. Patients and their family should be aware of the ongoing costs involved in optimising fertility. Fertility management may apply in both men and women. Fertility preservation options are different for men and women and the need for ongoing contraception applies to both men and women.

The potential for impaired fertility should be discussed and reinforced at different time points as appropriate throughout the diagnosis, treatment, surveillance and survivorship phases of care. These ongoing discussions will enable the patient and, if applicable, the family to make informed decisions. All discussions should be documented in the patient's medical record.



#### More information

Refer to the Cancer Council website <[www.cancervic.org.au/living-with-cancer/common-side-effects/fertility/fertility-overview.html](http://www.cancervic.org.au/living-with-cancer/common-side-effects/fertility/fertility-overview.html)> for more information.

### 3.6.3 Supportive care

Refer to the validated screening tools mentioned in Principle 4 'Supportive care'.

A number of specific challenges and needs may arise for patients at this time:

- assistance for dealing with psychological and emotional distress while adjusting to the diagnosis; treatment phobias; existential concerns; stress; difficulties making treatment decisions; anxiety or depression or both; psychosexual issues such as potential loss of fertility and premature menopause; history of sexual abuse; and interpersonal problems
- management of physical symptoms such as pain and fatigue (Australian Adult Cancer Pain Management Guideline Working Party 2019), nausea, vomiting, poor appetite, weight loss and itch
- malnutrition or undernutrition, identified using a validated nutrition screening tool such as the MST (note that many patients with a high BMI [obese patients] may also be malnourished [WHO 2018])
- nutritional support if weight loss is significant – refer to an accredited dietitian for assessment and support
- support for families or carers who are distressed with the patient's cancer diagnosis
- financial and employment issues (e.g. loss of income, travel and accommodation requirements for rural patients, and caring arrangements for other family members)
- support for families/relatives who may be distressed after learning of a genetically linked cancer diagnosis including counselling to support people to understand and adapt to the medical, psychological, familial and reproductive implications of their diagnosis
- specific spiritual needs that may benefit from the involvement of pastoral/spiritual care.

Additionally, palliative care may be required at this stage.

For more information on supportive care and needs that may arise for different population groups, refer to Appendices A, B and C.

### 3.6.4 Communication with patients, carers and families

In discussion with the patient, the lead clinician should undertake the following:

- establish if the patient has a regular or preferred general practitioner and if the patient does not have one, then encourage them to find one
- provide written information appropriate to the health literacy of the patient about the diagnosis and treatment to the patient and carer and refer the patient to the *Guide to best cancer care* (consumer optimal care pathway) for biliary cancer, as well as to relevant websites and support groups as appropriate
- provide a treatment care plan including contact details for the treating team and information on when to call the hospital
- discuss a timeframe for diagnosis and treatment with the patient and carer
- discuss the benefits of multidisciplinary care and gain the patient's consent before presenting their case at an MDM
- provide brief advice and refer to Quitline (13 7848) for behavioural intervention if the patient currently smokes (or has recently quit), and prescribe smoking cessation pharmacotherapy, if clinically appropriate

- recommend an 'integrated approach' throughout treatment regarding nutrition, exercise and minimal or no alcohol consumption among other considerations
- communicate the benefits of continued engagement with primary care during treatment for managing comorbid disease, health promotion, care coordination and holistic care
- where appropriate, review fertility needs with the patient and refer for specialist fertility management (including fertility preservation, contraception, management during pregnancy and of future pregnancies)
- be open to and encourage discussion about the diagnosis, prognosis (if the patient wishes to know) and survivorship and palliative care while clarifying the patient's preferences and needs, personal and cultural beliefs and expectations, and their ability to comprehend the communication
- while accepting that this is voluntary, encourage the patient to participate in advance care planning including considering appointing one or more substitute decision-makers and completing an advance care directive to clearly document their treatment preferences. Each state and territory has different terminology and legislation surrounding advance care directives and substitute decision-makers.

### 3.6.5 Communication with the general practitioner

The lead clinician has these communication responsibilities:

- involving the general practitioner from the point of diagnosis
- ensuring regular and timely communication with the general practitioner about the diagnosis, treatment plan and recommendations from MDMs and inviting them to participate in MDMs (consider using virtual mechanisms)
- supporting the role of general practice both during and after treatment
- discussing shared or team care arrangements with general practitioners or regional cancer specialists, or both, together with the patient.



#### More information

Refer to Principle 6 'Communication' for communication skills training programs and resources.

## Step 4: Treatment

**Step 4 describes the optimal treatments for biliary cancer, the training and experience required of the treating clinicians and the health service characteristics required for optimal cancer care.**

All health services must have clinical governance systems that meet the following integral requirements:

- identifying safety and quality measures
- monitoring and reporting on performance and outcomes
- identifying areas for improvement in safety and quality (ACSQHC 2020).

Step 4 outlines the treatment options for biliary cancer. For detailed clinical information on treatment options refer to these resources (preference is for more recent guidelines):

1. Clinical practice guideline interim update on the management of biliary tract cancer ESMO (2025) <[https://www.esmooopen.com/article/S2059-7029\(24\)01773-3/fulltext](https://www.esmooopen.com/article/S2059-7029(24)01773-3/fulltext)>
2. International Delphi consensus recommendations for gallbladder cancer, clinical practice guidelines. IHPBA-APHPBA (2025) <[https://www.hpbonline.org/article/S1365-182X\(24\)02223-8/abstract](https://www.hpbonline.org/article/S1365-182X(24)02223-8/abstract)>
3. Biliary tract cancers: French national clinical practice guidelines for diagnosis, treatments and follow-up (TNCD, SNFGE, FFCD, UNICANCER, GERCOR, SFCD, SFED, AFEF, SFRO, SFP, SFR, ACABi, ACHBPT) (2024) <[https://www.ejancer.com/article/S0959-8049\(24\)00156-4/abstract](https://www.ejancer.com/article/S0959-8049(24)00156-4/abstract)>
4. Practice guidelines for managing extrahepatic biliary tract cancers: The Korean Society of Hepato-Biliary-Pancreatic Surgery in collaboration with related societies (Korean Pancreatic and Biliary Surgery Society, Korean Society of Abdominal Radiology, Korean Society of Medical Oncology, Korean Society of Radiation Oncology, Korean Society of Pathologists, and Korean Society of Nuclear Medicine) (2024) <<https://www.ahbps.org/journal/view.html?doi=10.14701/ahbps.23-170>>
5. A practical guide for the systemic treatment of biliary tract cancer in Canada (2024) <[https://www.cholangio.ca/\\_files/ugd/8ff0b9\\_ab7a2e3619914e4d92f81f2bca9ed041.pdf](https://www.cholangio.ca/_files/ugd/8ff0b9_ab7a2e3619914e4d92f81f2bca9ed041.pdf)>
6. Transplantation Society of Australia and New Zealand (TSANZ) Clinical guidelines for organ transplantation from deceased donors, Version 1.13 (2024) <[https://tsanz.com.au/storage/Guidelines/TSANZ\\_Clinical\\_Guidelines\\_Version-113\\_FINAL.pdf](https://tsanz.com.au/storage/Guidelines/TSANZ_Clinical_Guidelines_Version-113_FINAL.pdf)>
7. Clinical practice guidelines on the management of intrahepatic cholangiocarcinoma. EASL-ILCA (2023) <[https://www.journal-of-hepatology.eu/article/S0168-8278\(23\)00185-X/fulltext](https://www.journal-of-hepatology.eu/article/S0168-8278(23)00185-X/fulltext)>
8. Biliary tract cancer: The European Society of Medical Oncology (ESMO) clinical practice guidelines for diagnosis, treatment, and follow-Up (2022) <<https://www.esmo.org/guidelines/esmo-clinical-practice-guideline-biliary-tract-cancer>>
9. Clinical practice guidelines for the management of biliary tract cancers 2019: the 3rd English edition. Japanese Society of Hepato-Biliary-Pancreatic Surgery <<https://onlinelibrary.wiley.com/doi/full/10.1002/jhbp.870>>
10. Adjuvant Therapy for Resected Biliary Tract Cancer: American Society of Clinical Oncology (ASCO) clinical practice guidelines (2018) <<https://www.ahpba.org/wp-content/uploads/2023/01/ASCO-adjuvant-therapy-resected-bile-tract-cancer.pdf>>

11. American International Hepato-Pancreato-Biliary Association (AHPBA) Intrahepatic Cholangiocarcinoma Consensus Statements (2014) <<https://pubmed.ncbi.nlm.nih.gov/26172134/>>
12. American International Hepato-Pancreato-Biliary Association (AHPBA) Hilar Cholangiocarcinoma Consensus Statements (2014) <<https://pubmed.ncbi.nlm.nih.gov/26172136/>>
13. American International Hepato-Pancreato-Biliary Association (AHPBA) Gallbladder Cancer Consensus Statements (2014) <<https://onlinelibrary.wiley.com/doi/abs/10.1111/hpb.12444>>.

#### 4.1 Treatment intent

The intent of treatment can be defined as one of the following:

- curative
- anti-cancer therapy to improve quality of life and/or longevity without expectation of cure
- symptom palliation.

The treatment intent should be established in a multidisciplinary setting, documented in the patient's medical record and conveyed to the patient and carer as appropriate.

The potential benefits need to be balanced against the morbidity and risks of treatment.

Patient comorbidities, including any co-existing liver disease, may affect treatment options and need to be treated appropriately. Optimisation of liver function (particularly addressing elevated bilirubin levels), plus other comorbidities, is usually required before and during treatment.

The lead clinician should discuss the advantages and disadvantages of each treatment and associated potential side effects with the patient and their carer or family before treatment consent is obtained and begins so the patient can make an informed decision. Supportive care services should also be considered during this decision-making process. Patients should be asked about their use of (current or intended) complementary therapies (refer to Appendix D).

A treatment plan may also be informed by genomic testing. Genomics-informed cancer care can improve the experience of people affected by cancer by optimising the treatment response and minimising treatment toxicity.

Timeframes for starting treatment should be informed by evidence-based guidelines where they exist. The treatment team should recognise that shorter timeframes for appropriate consultations and treatment can promote a better experience for patients.

Initiate advance care planning discussions with patients before treatment begins (this could include appointing a substitute decision-maker and completing an advance care directive). Formally involving a palliative care team/service may benefit any patient, so it is important to know and respect each person's preference.

## 4.2 Treatment options

### 4.2.1 Surgery

The potential for curative surgery depends on the staging of the tumour. Only a small percentage of patients have clearly resectable disease after careful clinical staging, due to advanced disease, frailty or comorbidities. This potential is assessed by the multidisciplinary team. Surgical resection offers the best long-term survival chance in patients with localised biliary cancer, although often this involves other treatments (e.g. chemotherapy or radiation treatment) as part of the surgical approach. Palliative surgery for metastatic cancer is not usually recommended.

#### Intrahepatic biliary cancer

- For intrahepatic biliary cancer, a partial hepatectomy (liver resection) may be performed including resection of the involved intra and extrahepatic bile ducts. Lymphadenectomy of regional nodes is recommended and includes an intraoperative assessment of any suspicious regional nodes, and ideally the recovery of at least six nodes.
- Portal vein embolisation is a safe and effective strategy for increasing the future liver remnant (FLR) before resection and may be most useful in patients with an FLR of less than 40 per cent.
- In the case of intrahepatic biliary cancer, liver transplantation may be considered for small lesions. Refer to the section below for recommendations for liver transplantation.

#### Perihilar biliary cancers

- For perihilar biliary cancers, either a right extended hepatectomy or left hemi-hepatectomy with concomitant bile duct resection and porta hepatis lymphadenectomy with bilioenteric reconstruction may be appropriate. For tumours with extensive bile duct involvement, a combined pancreaticoduodenectomy and hepatectomy may be needed to achieve negative margins.
- Portal vein embolisation is a safe and effective strategy for increasing the future liver remnant before resection and may be most useful in patients with an FLR of less than 40 per cent.
- Those with confirmed, or who are highly suspected of having, locally unresectable perihilar biliary cancer (confirmed by two HPB surgeons after discussion at an MDM) may also be considered for liver transplantation. Refer to the section below for recommendations for liver transplantation.

#### Distal biliary cancer

- Surgical resection of distal biliary cancers may be achieved with a pancreaticoduodenectomy, also known as Whipple's procedure. Further information on staging and treatment of distal biliary cancers requiring pancreaticoduodenectomy is in the optimal care pathway for people with pancreatic cancer.

#### Gallbladder cancer

- Surgery to remove the gallbladder is cholecystectomy. For gallbladder cancer, in addition to the gallbladder being removed, the adjacent liver portion, regional lymph nodes, and sometimes the bile duct, may need to be removed.
- Often gallbladder cancer is found incidentally on pathological examination after a simple cholecystectomy for cholecystitis or cholelithiasis. Patients with incidentally identified T1b, T2 or T3 disease in a cholecystectomy specimen should undergo re-resection with adequate lymphadenectomy, including an intraoperative assessment of any suspicious regional nodes, and ideally the recovery of at least six nodes, unless this is contraindicated by advanced disease or poor performance status.

### Special consideration: liver transplantation

- In the case of intrahepatic biliary cancer, liver transplantation may be considered for lesions smaller than 2 cm. For perihilar biliary cancer, a liver transplant may be considered for lesions no larger than 3 cm (TSANZ 2024). For both sites of biliary cancer, there needs to be clear evidence that there are no metastases including lymph node and peritoneal metastasis to consider a liver transplant.
- The diagnostic and treatment planning phase of a patient who may be considered for a liver transplant is highly specific. If the treatment team suspects a liver transplant may be an option, then referral and discussion through the MDM that includes a liver transplant surgeon is required as early as possible. Consideration is required in such instances to ensure the careful planning of biopsy, stenting and the role of neoadjuvant chemotherapy and/or radiotherapy.

### Timeframe for starting treatment

Surgery should be undertaken **within four weeks** of the MDM, depending on urgency, and when appropriate to the overall treatment plan.

### Training and experience required of the surgeon

Fellow of the Royal Australian College of Surgeons (or equivalent) with adequate training and experience that enables institutional credentialing and agreed scope of practice in HPB cancer and/or liver transplantation if relevant.

Documented evidence of the surgeon's training and experience, including their specific (sub-specialty) experience with hepatobiliary cancer and procedures to be undertaken, should be available.

International evidence suggests that surgeons who undertake a higher volume of resections have better clinical outcomes for complex cancer surgery such as HPB resections, although institutional volume is likely to be a more important factor than the individual surgeon (Gruen et al. 2009; Killeen et al. 2005). Patients undergoing HPB cancer surgery should be treated at specialist centres that have teams that deliver integrated expertise in endoscopy, imaging, interventional radiology, surgery and histopathology, and treat a high volume of these cases. Centres that do not meet these criteria should routinely refer cases to a centre with experience in that type of case.

### Health service characteristics

To provide safe and quality care for patients having surgery, health services should have these features:

- appropriate ward staff including nursing, dietetics, physiotherapy, occupational therapy and theatre resources to manage complex surgery
- critical care support
- 24-hour medical staff availability
- 24-hour operating room access
- intensive care unit with nursing and medical staff who are familiar with HPB surgery
- diagnostic imaging and interventional radiology
- ERCP
- access to endoscopic ultrasound
- pathology
- nuclear medicine imaging
- 24-hour access to interventional radiology fully supported by other surgical specialties
- specialist anaesthetists
- advanced endoscopy services
- full support from other surgical specialties.

Each unit performing complex HPB cancer surgery must demonstrate robust audit processes, and benchmarking of outcomes, to ensure quality outcomes are delivered consistently.

#### 4.2.2 Locoregional therapy

The choice of locoregional therapies is case specific, with decisions based on patient and tumour factors. It also depends on the outcomes of the MDM and the availability of local expertise in the treating centre.

### Radiation therapy

- External beam radiation therapy (EBRT), chemoradiation (CRT) or stereotactic body radiotherapy or stereotactic ablative radiotherapy (SBRT/SABR) may be appropriate.
- Radiotherapy may be considered for patients with:
  - unresectable or locally advanced biliary cancer, after systemic therapy is complete and the disease stabilised
  - intrahepatic biliary cancer with or without vascular involvement for patients in whom surgery is contraindicated
  - metastatic intrahepatic biliary cancer to reduce the risk of liver failure from local disease progression.
- Adjuvant CRT may be considered after margin positive resection (R1) or resection of node-positive disease.
- Adjuvant CRT may be considered following R0 resection of T2 or N1 gallbladder cancer.

### Intra-arterial treatment

- May be either transarterial chemoembolisation (TACE), selective internal radiation therapy (SIRT) or hepatic arterial infusion chemotherapy (HAIC).

### Thermal ablation

- Radiofrequency ablation (RFA) or microwave ablation (MWA) may be appropriate.
- Consider this for patients with intrahepatic biliary cancer no greater than 3 cm who have contraindications to surgery.

### Timeframe for starting treatment

Radiation therapy should begin **within four weeks** of the MDM, depending on urgency and modality. Adjuvant radiotherapy in postoperative care should begin within eight weeks of surgery, depending on the patient's recovery.

### Training and experience required of the appropriate specialists

Fellow of the Royal Australian and New Zealand College of Radiologists (or equivalent) with adequate training and experience that enables institutional credentialing and agreed scope of practice in biliary cancer.

The training and experience of radiation oncologist should be documented.

### Health service unit characteristics

To provide safe and quality care for patients having radiation therapy, health services should have these features:

- linear accelerator (LINAC) capable of image-guided radiation therapy (IGRT)
- dedicated CT planning
- access to MRI and PET imaging
- automatic record-verify of all radiation treatments delivered
- a treatment planning system
- trained medical physicists, radiation therapists and nurses with radiation therapy experience
- coordination for combined therapy with systemic therapy, especially where facilities are not co-located
- participation in Australian Clinical Dosimetry Service audits
- an incident management system linked with a quality management system.

### 4.2.3 Systemic therapy

A number of patients may benefit from systemic therapy:

- Adjuvant chemotherapy is recommended for most patients following resection for biliary cancer, depending on their recovery.
- Palliative chemotherapy is recommended for most patients with advanced biliary cancer, unless contraindicated by frailty or comorbidities.
- There are several chemotherapy regimens shown to have a role in biliary cancer. These regimens now include immune checkpoint inhibitors.
- Loco-regional treatment options such as embolisation, chemoembolisation, SIRT therapy and HAIC therapy can be considered as stand-alone options or in combination with systemic therapy, under MDM guidance.
- Neoadjuvant or perioperative chemotherapy in biliary cancer may be considered for borderline resectable disease, or as part of a liver transplant protocol or clinical trial.

### Timeframes for starting treatment

Systemic therapy for unresectable biliary cancer should begin **within four weeks** of the MDM, depending on urgency and modality. Adjuvant chemotherapy in postoperative care should begin **within eight weeks** of surgery, depending on the patient's recovery.

### Training and experience required of the appropriate specialists

Medical oncologists must have training and experience of this standard:

- Fellow of the Royal Australian College of Physicians (or equivalent)
- adequate training and experience that enables institutional credentialing and agreed scope of practice within this area (ACSQHC 2015).

Cancer nurses should have accredited training in these areas:

- anti-cancer treatment administration
- specialised nursing care for patients undergoing cancer treatments, including side effects and symptom management
- the handling and disposal of cytotoxic waste (ACSQHC 2020).

Systemic therapy should be prepared by a pharmacist whose background includes this experience:

- adequate training in systemic therapy medication, including dosing calculations according to protocols, formulations and/or preparation.

In a setting where no medical oncologist is locally available (e.g. regional or remote areas), some components of less complex therapies may be delivered by a general practitioner or nurse with training and experience that enables credentialing and agreed scope of practice within this area. This should be in accordance with a detailed treatment plan or agreed protocol, and with communication as agreed with the medical oncologist or as clinically required.

The training and experience of the appropriate specialist should be documented.

### Health service characteristics

To provide safe and quality care for patients having systemic therapy, health services should have these features:

- a clearly defined path to emergency care and advice after hours
- access to diagnostic pathology including basic haematology and biochemistry, and imaging
- cytotoxic drugs prepared in a pharmacy with appropriate facilities
- occupational health and safety guidelines regarding handling of cytotoxic drugs, including preparation, waste procedures and spill kits (eviQ 2022)
- guidelines and protocols to deliver treatment safely (including dealing with extravasation of drugs)
- coordination for combined therapy with radiation therapy, especially where facilities are not co-located
- appropriate molecular pathology access.

#### 4.2.4 Targeted therapies and immunotherapy

The role of targeted therapies and immunotherapies is evolving rapidly. There are several approved targeted therapies available to treat patients with locally advanced, unresectable or metastatic biliary cancer, with many clinical trials ongoing. Molecular profiling is advised for all patients to guide appropriate targeted and immunotherapy choice. Testing of multiple genes is preferred. Encourage patients to take in clinical trials wherever possible.

#### 4.2.5 Emerging therapies

The key principle for precision medicine is prompt and clinically oriented communication and coordination with an accredited laboratory and pathologist. Tissue analysis is integral for access to emerging therapies and, as such, tissue specimens should be treated carefully to avoid wastage and preserve DNA/RNA quality and quantity and to enable additional histopathological or molecular diagnostic tests in certain scenarios.

### 4.3 Palliative care

Early referral to palliative care can improve the quality of life for people with cancer and in some cases may be associated with survival benefits (Haines 2011; Temel et al. 2010; Zimmermann et al. 2014). This is particularly true for cancers with poor prognosis.

The lead clinician should ensure patients receive timely and appropriate referral to palliative care services. Referral should be based on need rather than prognosis. Emphasise the value of palliative care in improving symptom management and quality of life to patients and their carers.

The 'Dying to Talk' resource may help health professionals when initiating discussions with patients about future care needs (refer to 'More information'). Ensure that carers and families receive information, support and guidance about their role in palliative care (Palliative Care Australia 2024).

Patients, with support from their family or carer and treating team, should be encouraged to consider appointing a substitute decision-maker and to complete an advance care directive.

Refer to Step 6 for a more detailed description of managing patients with recurrent, residual or metastatic disease.



#### More information

These online resources are useful:

- Advance Care Planning Australia <[www.advancedcareplanning.org.au](http://www.advancedcareplanning.org.au)>
- Care Search <<https://www.caresearch.com.au/>>
- Dying to Talk <[www.dyingtotalk.org.au](http://www.dyingtotalk.org.au)>
- Palliative Care resource kit <[www.health.gov.au/health-topics/palliative-care](http://www.health.gov.au/health-topics/palliative-care)>
- Palliative Care Australia (for patients and carers) <[www.palliativecare.org.au](http://www.palliativecare.org.au)>.

### 4.4 Research and clinical trials

The team should support the patient to take part in research or clinical trials where available and appropriate. Many emerging treatments are only available on clinical trials that may require referral to certain trial centres.

For more information visit the Cancer Australia website <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>.

## 4.5 Support and communication

### 4.5.1 Supportive care

Refer to the validated screening tools mentioned in Principle 4 'Supportive care'.

A number of specific challenges and needs may arise for patients at this time:

- assistance for dealing with emotional and psychological issues, including body image concerns, fatigue, quitting smoking, traumatic experiences, existential anxiety, treatment phobias, anxiety/depression, interpersonal problems and sexuality concerns
- potential isolation from normal support networks, particularly for rural patients who are staying away from home for treatment
- management of physical symptoms such as pain, fatigue, nausea, vomiting, poor appetite, weight loss and itch
- surgical procedures such as stent placement or biliary bypass to relieve symptoms of jaundice (when endoscopic access is not possible for stent placement, percutaneous transhepatic drainage is recommended; in patients with a life expectancy of more than three months, a metal stent is preferred)
- coeliac plexus neurolysis to prevent and/or manage intractable abdominal pain –abdominal pain is a common feature that may be due to local and/or coeliac plexus involvement, so patients with severe pain or pain requiring rapid escalation of opioids should be offered the procedure (coeliac plexus neurolysis should be undertaken in a centre that has collaboration with anaesthetists, interventional gastroenterologists, interventional radiologists, oncologists and palliative care physicians)
- dietary issues including malnutrition from disease or treatments resulting in need for referral to dietitian
- decline in mobility or functional status as a result of treatment and disease
- assistance with beginning or resuming regular exercise with referral to an exercise physiologist or physiotherapist (COSA 2018; Hayes et al. 2019)
- Early involvement of general practitioners may lead to improved cancer survivorship care following acute treatment. General practitioners can address many supportive care needs through good communication and clear guidance from the specialist team (Emery 2014).
- Patients, carers and families may have these additional issues and needs:
- financial issues related to loss of income (through reduced capacity to work or loss of work) and additional expenses as a result of illness or treatment
- advance care planning, which may involve appointing a substitute decision-maker and completing an advance care directive
- legal issues (completing a will, care of dependent children) or making an insurance, superannuation or social security claim on the basis of terminal illness or permanent disability.

Cancer Council's 13 11 20 information and support line can assist with information and referral to local support services.

For more information on supportive care and needs that may arise for different population groups, refer to Appendices A, B and C.

#### **4.5.2 Rehabilitation**

Rehabilitation may be required at any point of the care pathway. If it is required before treatment, it is referred to as prehabilitation (refer to section 3.6.1).

All members of the multidisciplinary team have an important role in promoting rehabilitation. Team members may include occupational therapists, speech pathologists, dietitians, social workers, psychologists, physiotherapists, exercise physiologists and rehabilitation specialists.

To maximise the safety and therapeutic effect of exercise for people with cancer, all team members should recommend that people with cancer work towards achieving, and then maintaining, recommended levels of exercise and physical activity as per relevant guidelines. Exercise should be prescribed and delivered under the direction of an accredited exercise physiologist or physiotherapist with experience in cancer care (Vardy et al. 2019). The focus of intervention from these health professionals is tailoring evidence-based exercise recommendations to the individual patient's needs and abilities, with a focus on the patient transitioning to ongoing self-managed exercise.

Other issues that may need to be dealt with include managing cancer-related fatigue, improving physical endurance, achieving independence in daily tasks, optimising nutritional intake, returning to work and ongoing adjustment to cancer and its sequels. Referrals to dietitians, psychosocial support, return-to-work programs and community support organisations can help in managing these issues.

#### **4.5.3 Communication with patients, carers and families**

The lead or nominated clinician should take responsibility for these tasks:

- discussing treatment options with patients and carers, including the treatment intent and expected outcomes, and providing a written version of the plan and any referrals
- providing patients and carers with information about the possible side effects of treatment, managing symptoms between active treatments, how to access care, self-management strategies and emergency contacts
- encouraging patients to use question prompt lists and audio recordings, and to have a support person present to aid informed decision-making
- initiating a discussion about advance care planning and involving carers or family if the patient wishes.

#### 4.5.4 Communication with the general practitioner

The general practitioner plays an important role in coordinating care for patients, including helping to manage side effects and other comorbidities, and offering support when patients have questions or worries. For most patients, simultaneous care provided by their general practitioner is very important.

The lead clinician, in discussion with the patient's general practitioner, should consider these points:

- the general practitioner's role in symptom management, supportive care and referral to local services
- using a chronic disease management plan and mental health care management plan
- how to ensure regular and timely two-way communication about:
  - the treatment plan, including intent and potential side effects
  - supportive and palliative care requirements
  - the patient's prognosis and their understanding of this
  - enrolment in research or clinical trials
  - changes in treatment or medications
  - the presence of an advance care directive or appointment of a substitute decision-maker
  - recommendations from the multidisciplinary team.



#### **More information**

Refer to Principle 6 'Communication' for communication skills training programs and resources.

## Step 5: Care after initial treatment and recovery

**The term ‘cancer survivor’ describes a person living with cancer, from the point of diagnosis until the end of life. Survivorship care in Australia has traditionally been provided to patients who have completed active treatment and are in the post-treatment phase. But there is now a shift to provide survivorship care and services from the point of diagnosis to improve cancer-related outcomes.**

Cancer survivors may experience inferior quality of life and cancer-related symptoms for up to five years after their diagnosis (Jefford et al. 2017). Distress, fear of cancer recurrence, fatigue, obesity and sedentary lifestyle are common symptoms reported by cancer survivors (Vardy et al. 2019).

Due to an ageing population and improvements in treatments and supportive care, the number of people surviving cancer is increasing. International research shows there is an important need to focus on helping cancer survivors cope with life beyond their acute treatment. Cancer survivors often face issues that are different from those experienced during active treatment for cancer and may include a range of issues, as well as unmet needs that affect their quality of life (Lisy et al. 2019; Tan et al. 2019).

Physical, emotional and psychological issues include fear of cancer recurrence, cancer-related fatigue, pain, distress, anxiety, depression, cognitive changes and sleep issues (Lisy et al. 2019). Late effects may occur months or years later and depend on the type of cancer treatment. Survivors and their carers may experience impacted relationships and practical issues including difficulties with return to work or study and financial hardship. They may also experience changes to sex and intimacy. Fertility, contraception and pregnancy care after treatment may require specialist input.

The Institute of Medicine, in its report *From cancer patient to cancer survivor: Lost in transition*, describes the essential components of survivorship care listed as:

- Prevention of recurrent and new cancers, and of other late effects
- Surveillance for cancer spread, recurrence or second cancers; assessment of medical and psychosocial late effects
- Intervention for consequences of cancer and its treatment, for example: medical problems such as lymphedema and sexual dysfunction; symptoms, including pain and fatigue; psychological distress experienced by cancer survivors and their caregivers; and concerns related to employment, insurance and disability
- Coordination between specialists and primary care providers to ensure that all the survivor’s health needs are met (Hewitt et al. 2006).

Access to a range of health professions may be required including physiotherapy, occupational therapy, social work, dietetics, clinical psychology, fertility and palliative care. Coordinating care between all providers is essential to ensure the patient’s needs are met.

Cancer survivors are more likely than the general population to have and/or develop comorbidities (Vijayvergia & Denlinger 2015). Health professionals should support survivors to self-manage their own health needs and to make informed decisions about lifestyle behaviours that promote wellness and improve their quality of life (Australian Cancer Survivorship Centre 2024; Cancer Australia 2017).

## 5.1 Transitioning from active treatment

The transition from active treatment to post-treatment care is critical to long-term health. In some cases, people will need ongoing, hospital-based care, and in other cases a shared follow-up care arrangement with their general practitioner may be appropriate. This will vary depending on the type and stage of cancer and needs to be planned.

Shared follow-up care involves the joint participation of specialists and general practitioners in the planned delivery of follow-up and survivorship care. A shared care plan is developed that outlines the responsibilities of members of the care team, the follow-up schedule, triggers for review, plans for rapid access into each setting and agreement regarding format, frequency and triggers for communication.

After completing initial treatment, a designated member of the multidisciplinary team (most commonly nursing or medical staff involved in the patient's care) should provide the patient with a needs assessment and treatment summary and develop a survivorship care plan in conjunction with the patient. This should include a comprehensive list of issues identified by all members of the multidisciplinary team involved in the patient's care and by the patient. These documents are key resources for the patient and their healthcare providers and can be used to improve communication and care coordination.

The treatment summary should cover, but is not limited to:

- the diagnostic tests performed and results
- diagnosis including stage, prognostic or severity score
- tumour characteristics
- treatment received (types and dates)
- current toxicities (severity, management and expected outcomes)
- interventions and treatment plans from other health providers
- potential long-term and late effects of treatment
- supportive care services provided
- follow-up schedule
- contact information for key healthcare providers.

## 5.2 Follow-up care

There is currently no high-level evidence to inform the frequency of clinical follow-up or the use of routine investigations for biliary cancer. Investigations should be directed to symptoms. Each centre should establish a regular surveillance protocol. Surveillance may consist of three- to six-monthly visits during the first two years and six- to 12-monthly visits for up to five years or as clinically indicated. A combination of clinical examination, laboratory investigation, tumour markers and an annual CT scan of the chest, abdomen and pelvis may be appropriate.

Responsibility for follow-up care should be agreed between the lead clinician, the general practitioner, relevant members of the multidisciplinary team and the patient. This is based on guideline recommendations for post-treatment care, as well as the patient's current and anticipated physical and emotional needs and preferences.

Evidence, in other cancer types, comparing shared follow-up care and specialised care indicates equivalence in outcomes including recurrence rate, cancer survival and quality of life (Cancer Research in Primary Care 2016).

Ongoing communication between healthcare providers involved in care and a clear understanding of roles and responsibilities is key to effective survivorship care.

In particular circumstances, other models of post-treatment care can be safely and effectively provided such as nurse-led models of care (Monterosso et al. 2019). Other models of post-treatment care can be provided in these locations or by these health professionals:

- in a shared care setting
- in a general practice setting
- by non-medical staff
- by allied health or nurses
- in a non-face-to-face setting (e.g. by telehealth).

A designated member of the team should document the agreed survivorship care plan. The survivorship care plan should support wellness and have a strong emphasis on healthy lifestyle changes such as avoiding tobacco, limiting or avoiding alcohol, a balanced diet, a non-sedentary lifestyle, weight management and a mix of aerobic and resistance exercise (COSA 2018; Hayes et al. 2019).

This survivorship care plan should also cover, but is not limited to:

- what medical follow-up is required (surveillance for recurrence or secondary and metachronous cancers, screening and assessment for medical and psychosocial effects)
- model of post-treatment care, the health professional providing care and where it will be delivered
- care plans from other health providers to manage the consequences of cancer and cancer treatment
- wellbeing, primary and secondary prevention health recommendations that align with chronic disease management principles
- rehabilitation recommendations
- available support services
- a process for rapid re-entry to specialist medical services for suspected recurrence.

Survivors generally need regular follow-up, often for five or more years after cancer treatment finishes. The survivorship care plan therefore may need to be updated to reflect changes in the patient's clinical and psychosocial status and needs.

Processes for rapid re-entry to hospital care should be documented and communicated to the patient and relevant stakeholders.

Care in the post-treatment phase is driven by predicted risks (e.g. the risk of recurrence, developing late effects of treatment and psychological issues) as well as individual clinical and supportive care needs. It is important that post-treatment care is based on evidence and is consistent with guidelines. Not all people will require ongoing tests or clinical review and may be discharged to general practice follow-up.

The lead clinician should discuss (and general practitioner reinforce) options for follow-up at the start and end of treatment. It is critical for optimal aftercare that the designated member of the treatment team educates the patient about the symptoms of recurrence.

General practitioners (including nurses) can:

- connect patients to local community services and programs
- manage long-term and late effects
- manage comorbidities
- provide wellbeing information and advice to promote self-management
- screen for cancer and non-cancerous conditions.



### More information

Templates and other resources to help with developing treatment summaries and survivorship care plans are available from these organisations:

- Australian Cancer Survivorship Centre
- Cancer Australia – Principles of Cancer Survivorship
- Cancer Council Australia and states and territories
- Clinical Oncology Society of Australia – Model of Survivorship Care
- eviQ – Cancer survivorship: introductory course
- mycareplan.org.au
- South Australian Cancer Service – Statewide Survivorship Framework resources
- American Society of Clinical Oncology – guidelines.

#### 5.2.1 Preventing recurrence

Not smoking, eating a healthy diet, being sun smart, avoiding or limiting alcohol intake, being physically active and maintaining a healthy body weight may help reduce the risk of primary recurrence or a second primary cancer.

Encourage and support all cancer survivors to reduce modifiable risk factors for recurrence as well as other chronic diseases. Ongoing coordination of care between providers should also deal with any comorbidities, particularly ongoing complex and life-threatening comorbid conditions.

#### 5.3 Research and clinical trials

Support cancer survivors to participate in research or clinical trials where they are available and appropriate. These might include studies to understand survivors' issues, to better manage treatment side effects, or to improve models of care and quality of life.

For more information visit the Cancer Australia website <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>.

## **5.4 Support and communication**

### **5.4.1 Supportive care**

Refer to the validated screening tools mentioned in Principle 4 'Supportive care'. The 'Cancer Survivors Unmet Needs (CaSun)' is another validated screening tool that may help health professionals to identify the unmet needs of patients during survivorship.

A number of specific challenges and needs may arise for cancer survivors:

- financial and employment issues (e.g. loss of income and assistance with returning to work, and the cost of treatment, travel and accommodation)
- appointing a substitute decision-maker and completing an advance care directive
- legal issues such as completing a will.

For more information on supportive care and needs that may arise for different population groups, refer to Appendices A, B and C.

### **5.4.2 Rehabilitation and recovery**

Rehabilitation may be required at any point of the care pathway from the pre-treatment phase through to disease-free survival and palliative care (Cormie et al. 2017).

Issues that may need to be dealt with include managing cancer-related fatigue, coping with cognitive changes, improving physical endurance, achieving independence in daily tasks, returning to study or work and ongoing adjustment to cancer and its sequels.

Exercise is a safe and effective intervention that improves the physical and emotional health and wellbeing of cancer patients. Exercise should be embedded as part of standard practice in cancer care and be viewed as an adjunct therapy that helps counteract the adverse effects of cancer and its treatment.

Cancer survivors may find referral to specific cancer rehabilitation, optimisation programs or community-based rehabilitation appropriate and beneficial. Other options include referral to allied health supports through team care arrangements and mental health plans. Some community support organisations (cancer-related non-government, not-for-profit and charities) provide services to cancer survivors.

### **5.4.3 Communication with patients, carers and families**

The lead clinician (themselves or by delegation) should take responsibility for these tasks:

- explaining the model of post-treatment care and the roles of health professionals involved in post-treatment care including the role of general practice
- explaining the treatment summary and follow-up care plan
- discussing the development of a shared follow-up and survivorship care plan where a model of shared follow-up care has been agreed
- discussing how to manage any of the physical, psychological or emotional issues identified
- providing information on the signs and symptoms of recurrent disease
- providing a survivorship care plan with information on secondary prevention and healthy living
- providing contact details of the care team involved
- providing clear information about the role and benefits of palliative care and advance care planning.

#### 5.4.4 Communication with the general practitioner

The lead clinician should ensure regular, timely, two-way communication with the general practitioner about:

- the patient's progress
- the follow-up care plan
- potential late effects
- supportive and palliative care requirements
- any shared care arrangements
- clarification of various roles in patient care
- a process for rapid re-entry to medical services for patients with suspected recurrence or if there are other concerns.



#### More information

Refer to Principle 6 'Communication' for communication skills training programs and resources.

## Step 6: Managing recurrent, residual or metastatic disease

**Patients who present with recurrent, residual or metastatic disease should be managed by a multidisciplinary team and offered timely referral to appropriate physical, practical and emotional support.**

Step 6 is concerned with managing recurrent or local residual and metastatic disease. The likelihood of recurrence depends on many factors usually related to the type of cancer, the stage of cancer at presentation and the effectiveness of treatment. Some cancers cannot be eradicated even with the best initial treatment. But controlling disease and disease-related symptoms is often possible, depending on the clinical situation.

### 6.1 Signs and symptoms of metastatic disease

Some patients will have metastatic disease on initial presentation. Others may present with symptoms of recurrent disease after a previous cancer diagnosis. Access to the best available therapies, including clinical trials, as well as treatment overseen by a multidisciplinary team, are crucial to achieving the best outcomes for anyone with metastatic disease.

Signs and symptoms will depend on the type of cancer initially diagnosed and the location of metastatic disease. They may be discovered by the patient or by surveillance in the post-treatment period. For biliary cancer, signs and symptoms of metastatic disease are often the same as for initial presentation. Refer to Step 2.

### 6.2 Managing metastatic disease

Managing metastatic disease is complex and should therefore involve all the appropriate specialties in a multidisciplinary team including palliative care where appropriate. Treatment of recurrent, residual or metastatic biliary cancer is rarely curative; metastatic biliary cancer usually has a poor prognosis. For most patients, treatment is given with palliative rather than curative intent.

From the time of diagnosis, the team should offer patients appropriate psychosocial care, supportive care, advance care planning and symptom-related interventions as part of their routine care. The approach should be personalised to meet the patient's individual needs, values and preferences. The full complement of supportive care measures as described throughout the optimal care pathway and in Appendices A, B and C should be offered to assist patients and their families and carers to cope. These measures should be updated as the patient's circumstances change.

Survivorship care should be considered and offered at an early stage. Many people live with advanced cancer for many months or years. As survival is improving in many patients, survivorship issues should be considered as part of routine care. Health professionals should therefore be ready to change and adapt treatment strategies according to disease status, prior treatment tolerance and toxicities and the patient's quality of life, in addition to the patient's priorities and life plans.

### 6.3 Multidisciplinary team

If there is an indication that a patient's cancer has returned, care should be provided under the guidance of a treating specialist. Each patient should be evaluated to determine if referral to the original multidisciplinary team is necessary. Often referral back to the original multidisciplinary team will not be necessary unless there are obvious aspects of care involving different therapeutic and supportive care disciplines not otherwise accessible. The multidisciplinary team may include new members such as palliative care specialists.

## 6.4 Treatment

Treatment will depend on the location, extent of recurrent or residual disease, previous management and the patient's preferences.

In managing people with recurrent, residual or metastatic biliary cancer, treatment may include these options:

- management of biliary sepsis – biliary obstruction and sepsis are an important source of morbidity that may need tailored care to manage; nevertheless, managing biliary sepsis is often urgent because these infections can be life-threatening and may require immediate supportive measures as well as rapid initiation of intravenous antibiotics (once samples for blood culture have been collected)
- surgical procedures such as stent placement or biliary bypass to relieve symptoms of jaundice (when endoscopic access is not possible, percutaneous transhepatic drainage is recommended; stent insertion and replacement may be required – in patients with a life expectancy of greater than three months, a metal stent is preferred)
- internal or external radiotherapy as palliative treatment to relieve symptoms such as pain or to shrink tumours that may be causing pain
- chemotherapy
- immunotherapy
- targeted therapy
- radiofrequency ablation (radiofrequency waves are delivered to unresectable tumours with the goal to thermally ablate the tumour – this is an option for patients with biliary cancer whose tumours are smaller than 3 cm but are not amenable to surgery)
- transarterial chemoembolisation
- pain management – consider coeliac plexus neurolysis to prevent and/or manage intractable abdominal pain; offer the procedure to patients with severe pain or pain requiring rapid escalation of opioids (coeliac plexus neurolysis should be undertaken in a centre that has collaboration with anaesthetists, interventional gastroenterologists, interventional radiologists, oncologists and palliative care physicians)
- palliative care.

Treatment may focus on disease control or palliation, based on the extent of the disease, general health or the patient's preferences and values. The potential goals of treatment should be discussed, respecting the patient's cultural values. Wherever possible, written information should be provided. Encourage early referral to clinical trials or accepting an invitation to participate in research.

## 6.5 Advance care planning

Advance care planning is important for all patients with a cancer diagnosis to consider, but especially those with advanced disease. Patients should be encouraged to think and talk about their healthcare values and preferences with family or carers, appoint a substitute decision-maker and consider developing an advance care directive to convey their preferences for future health care in the event they become unable to communicate their wishes.



### More information

Refer to section 4.3 'More information' for links to resources.

Refer patients and carers to Advance Care Planning Australia <[www.advancecareplanning.org.au](http://www.advancecareplanning.org.au)> or to the Advance Care Planning National Phone Advisory Service on 1300 208 582.

## 6.6 Palliative care

Early referral to palliative care can improve the quality of life for people with cancer and in some cases may be associated with survival benefits (Haines 2011; Temel et al. 2010; Zimmermann et al. 2014). The treatment team should emphasise the value of palliative care in improving symptom management and quality of life to patients and their carers. Refer to section 4.3 for more detailed information.

The lead clinician should ensure timely and appropriate referral to palliative care services. Referral to palliative care services should be based on the patient's need and potential for benefit, not prognosis.



### More information

Refer to the end of section 4.3 'Palliative care' for links to resources.

## 6.7 Research and clinical trials

The treatment team should support the patient to participate in research and clinical trials where available and appropriate.

For more information visit the Cancer Australia website <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>.

## 6.8 Support and communication

### 6.8.1 Supportive care

Refer to the validated screening tools mentioned in Principle 4 'Supportive care'.

A number of specific challenges and needs may arise at this time for patients:

- assistance for dealing with emotional and psychological distress resulting from fear of death or dying, existential concerns, anticipatory grief, communicating wishes to loved ones, interpersonal problems and sexuality concerns
- potential isolation from normal support networks, particularly for rural patients who are staying away from home for treatment
- cognitive changes as a result of treatment and disease progression such as altered memory, attention and concentration (a patient may appoint someone to make medical, financial and legal decisions on their behalf – a substitute decision-maker – before and in case they experience cognitive decline)
- management of physical symptoms including pain, fatigue, nausea, vomiting, poor appetite, weight loss and itch
- dietary issues including malnutrition as a result of disease or treatments (referral to dietitian may be required)
- decline in mobility or functional status as a result of recurrent disease and treatments (referral to physiotherapy or occupational therapy may be required)
- coping with hair loss and changes in physical appearance (refer to the *Look Good, Feel Better* program – refer to 'Resource list')
- appointing a substitute decision-maker and completing an advance care directive
- financial issues as a result of disease recurrence such as gaining early access to superannuation and insurance
- legal issues (completing a will, care of dependent children) and making an insurance, superannuation or social security claim on the basis of terminal illness or permanent disability.

### 6.8.2 Rehabilitation

Rehabilitation may be required at any point of the metastatic care pathway, from preparing for treatment through to palliative care. Issues that may need to be dealt with include managing cancer-related fatigue, improving physical endurance, achieving independence in daily tasks, returning to work and ongoing adjustment to cancer and its sequels.

Exercise is a safe and effective intervention that improves the physical and emotional health and wellbeing of cancer patients. Exercise should be embedded as part of standard practice in cancer care and be viewed as an adjunct therapy that helps counteract the adverse effects of cancer and its treatment.

### 6.8.3 Communication with patients, carers and families

The lead clinician should ensure there is adequate discussion with patients and carers about the diagnosis and recommended treatment, including treatment intent and possible outcomes, likely adverse effects and the supportive care options available. The lead clinician should also be open to and encourage discussion about the prognosis (if the patient wishes to know), survivorship and palliative care while clarifying the patient's preferences and needs, personal and cultural beliefs and expectations, and their ability to comprehend the communication. Patients should be advised of the likely duration of stent patency and of symptoms and signs that indicate biliary obstruction or infection.



#### More information

Refer to Principle 6 'Communication' for communication skills training programs and resources.

## Step 7: End-of-life care

**Step 7 is concerned with maintaining the patient's quality of life and meeting their health and supportive care needs as they approach the end of life, as well as the needs of their family and carers.**

Some patients with advanced cancer will reach a time when active treatment is no longer appropriate. The team needs to share the principles of a palliative approach to care when making decisions with the patient and their family or carer. End-of-life care is appropriate when the patient's symptoms are increasing and functional status is declining.

### 7.1 Multidisciplinary palliative care

If the treatment team does not include a palliative care member, the lead clinician should consider referring the patient to palliative care services, with the general practitioner's engagement. This may include inpatient palliative unit access (as required).

The multidisciplinary team may consider seeking additional expertise from these professionals:

- clinical psychologist
- clinical nurse specialist or practitioner
- social worker
- palliative medicine specialist
- pain specialist
- pastoral or spiritual carer
- bereavement counsellor
- music therapist
- art therapist
- cultural expert
- Canteen for children of parents with cancer <<https://www.canteen.org.au/young-people/parent-carer-cancer>>.

The team might also recommend that patients access these services:

- home and community-based care
- specialist community palliative care workers
- community nursing.

If the patient does not already have an advance care directive in place, a designated member of the treatment team should ensure the patient has been informed about advance care directives and their benefits. They should be offered an opportunity to develop one in collaboration with their family or carer, if they choose.

The patient's wishes and preferences for their end-of-life care, including an advance care directive if the patient has one, should be shared with treating teams and should guide care decisions if the patient is not able to speak for themselves.

It is essential for the treatment team to consider the appropriate place of care, the patient's preferred place of death and the support needed for the patient, their family and carers.

The treatment team should also ensure that carers and families receive the information, support and guidance about their role according to their needs and wishes (Palliative Care Australia 2024).



### More information

The treatment team can refer patients and carers to these resources:

- Palliative Care Australia <[www.palliativecare.org.au](http://www.palliativecare.org.au)>
- Advance Care Planning Australia <[www.advancecareplanning.org.au](http://www.advancecareplanning.org.au)> or to Advance Care Planning Australia's National Advisory Service on 1300 208 582.

#### 7.1.1 Voluntary Assisted Dying

A person who is suffering in the late stages of advanced disease and who meets strict criteria can request access to voluntary assisted dying. It must be voluntary and initiated by the person themselves.

As voluntary assisted dying is legislated by state and territory governments, it is essential to know the rules around this choice in the state or territory where the patient lives (Cancer Council 2020). Contact your jurisdiction's health department for the latest voluntary assisted dying information relevant to your state/territory.



### More information

Contact your jurisdiction's health department <<https://www.health.gov.au/about-us/contact-us/state-and-territory-health-departments>> for the latest voluntary assisted dying information relevant to your state or territory.

## 7.2 Research and clinical trials

Clinical trials may help improve palliative care and in managing a patient's symptoms of advanced cancer (Cancer Council Victoria 2019). The treatment team should support the patient to participate in research and clinical trials where available and appropriate.

For more information visit the Cancer Australia website <[www.australiancancertrials.gov.au](http://www.australiancancertrials.gov.au)>. Refer to the 'Resource list' for additional clinical trial databases.

## 7.3 Support and communication

### 7.3.1 Supportive care

Refer to the validated screening tools mentioned in Principle 4 'Supportive care'.

A number of specific challenges and needs may arise for patients at this time:

- assistance for dealing with emotional and psychological distress from anticipatory grief, fear of death or dying, anxiety/depression and interpersonal problems
- management of physical symptoms associated with biliary obstruction such as pain, itch and cholangitis, as well as fatigue, poor appetite and weight loss, nausea and vomiting
- management of biliary sepsis – biliary obstruction and sepsis are an important source of morbidity that may require tailored care to manage; nevertheless, managing biliary sepsis is often urgent because these infections can be life-threatening and may require immediate supportive measures as well as rapid initiation of intravenous antibiotics (once samples for blood culture have been collected)
- biliary drainage and subsequent treatment in patients with biliary obstruction – when endoscopic access is not possible, percutaneous transhepatic drainage is recommended (stent insertion and replacement may be required; in patients with a life expectancy of more than three months, a metal stent is preferred)

- coeliac plexus neurolysis to manage intractable abdominal pain – abdominal pain is a common feature that may be due to local and/or coeliac plexus involvement so patients with severe pain or pain requiring rapid escalation of opioids should be offered the procedure (coeliac plexus neurolysis should be undertaken in a centre that has collaboration with anaesthetists, interventional gastroenterologists, interventional radiologists, oncologists and palliative care physicians)
- decline in mobility or functional status, affecting the patient's discharge destination (a referral to physiotherapy, exercise physiology, occupational therapy or social work may be needed)
- appointing a substitute decision-maker and completing an advance care directive
- legal issues (completing a will, care of dependent children) and making an insurance, superannuation or social security claim on the basis of terminal illness or permanent disability
- specific support for families where a parent is dying and will leave behind bereaved children or adolescents, creating special family needs
- arranging a funeral.
- These services and resources can help:
  - referral to 13 11 20 for Cancer Council Australia's Pro Bono Program for free legal, financial, small business accounting and workplace assistance (subject to a means test)
  - *Sad news sorry business* (Queensland Health 2015) for the specific needs of Aboriginal and Torres Strait Islander people.

For more information on supportive care and needs that may arise for different population groups, refer to Appendices A, B and C.

### 7.3.2 Communication with patients, carers and families

The lead clinician is responsible for:

- being open to and encouraging discussion with the patient about the expected disease course, considering the patient's personal and cultural beliefs and expectations
- advising patients of the likely duration of stent patency and of symptoms and signs that indicate biliary obstruction or infection
- discussing palliative care options, including inpatient and community-based services as well as dying at home and subsequent arrangements
- providing the patient and carer with the contact details of a palliative care service
- referring the patient to palliative care in the community according to the carer's wishes.

### 7.3.3 Communication with the general practitioner

The lead clinician should discuss end-of-life care planning to ensure the patient's needs and goals are met in the appropriate environment. The patient's general practitioner should be kept fully informed and involved in major developments in the patient's illness path.



#### More information

For support with communication skills and training programs, refer to these sources:

- *Sad news sorry business* <[www.health.qld.gov.au/\\_\\_data/assets/pdf\\_file/0023/151736/sorry\\_business.pdf](http://www.health.qld.gov.au/__data/assets/pdf_file/0023/151736/sorry_business.pdf)>
- Principle 6 'Communication'.

# Contributors and reviewers

Our thanks to the following health professionals, consumer representatives, stakeholders and organisations consulted in developing this optimal care pathway.

## Expert working group

Professor John Zalcborg AO (Chair), Medical Oncologist, Alfred Health and Monash University, Vic

Alana Fitzgibbon, Upper Gastrointestinal Cancer Clinical Nurse Consultant, Royal Hobart Hospital, Tas

Catherine Trevaskis, Gastrointestinal Cancer Specialist Nurse, Canberra Health Services, ACT

Associate Professor Charles Pilgrim, Surgeon, Alfred Health and Monash University, Vic

Professor David Goldstein, Medical Oncologist, Prince of Wales Hospital and University of NSW, NSW

Grant Rutley, Consumer and Founder Battle Upper GI Cancer, Vic

Dr Javiera Martinez Gutierrez, GP, Victorian Comprehensive Cancer Centre and University of Melbourne, Vic

Dr Julie Chu, Radiation Oncologist, Peter MacCallum Cancer Centre, Vic

Dr Julie Lokan, Pathologist, Austin Health, Vic

Professor Lorraine Chantrill, Medical Oncologist, Illawarra Shoalhaven Local Health District, NSW

Melanie Regan, Project Manager and Cancer Nurse, Pancare Foundation, Vic

Dr Nicola Atkin, Palliative Care Physician, Peter MacCallum Cancer Centre and The Royal Melbourne Hospital, Vic

Steve Holmes, Consumer and Founder Cholangiocarcinoma Foundation Australia, Qld

Professor Stewart Dunn, Psychologist, Royal North Shore Hospital and the University of Sydney, NSW

Tiffany Dobson, Upper Gastrointestinal Cancer Support Nurse, Pancare Foundation, Vic

And the 171 multidisciplinary clinicians from across Australia who contributed to this optimal care pathway via the national Biliary Tract Cancer Consensus Guidelines Group.

## Medical colleges and peak organisations invited to provide feedback

Advance Care Planning Australia

Allied Health Professions Australia

Australasian Association of Nuclear Medicine Specialists

Australasian Chapter of Palliative Medicine, Royal Australia College of Physicians

Australasian Gastro-Intestinal Trials Group

Australia and New Zealand Hepatic, Pancreatic and Biliary Association

Australian and New Zealand Society of Neuroradiology

Australian and New Zealand Society of Palliative Care

Australian Cancer Survivorship Centre

Australian College of Nursing

Australian Medical Association

Australian Society of Medical Imaging and Radiation Therapy

Cancer Nurses Society of Australia

Clinical Oncology Society of Australia

Dietitians Australia

Gastroenterological Nurses College of Australia

Gastroenterological Society of Australia

Interventional Radiology Society of Australasia

Liver Foundation

Medical Oncology Group of Australia

Oncology Social Workers Australia and New Zealand

Psycho-Oncology Co-operative Research Group

Royal Australasian College of Physicians

Royal Australasian College of Surgeons

Royal Australian and New Zealand College of Radiologists

Royal Australian College of General Practitioners

Royal College of Pathologists of Australasia

Upper Gastrointestinal Cancer Registry


# Appendix A: Supportive care domains

Supportive care in cancer refers to the following five domains:

- the physical domain, which includes a wide range of physical symptoms that may be acute, relatively short lived or ongoing, requiring continuing interventions or rehabilitation
- the psychological domain, which includes a range of issues related to the patient's mental health wellbeing and personal relationships
- the social domain, which includes a range of social and practical issues that will affect the patient, carer and family such as the need for emotional support, maintaining social networks and financial concerns
- the information domain, which includes access to information about cancer and its treatment, recovery and survivorship support services and the health system overall
- the spiritual domain, which focuses on the patient's changing sense of self and challenges to their underlying beliefs and existential concerns (Palliative Care Victoria 2025).

Fitch's (2000) model of supportive care recognises the variety and level of intervention required at each critical point as well as the need to be specific to the individual patient (Figure A1). The model targets the type and level of intervention required to meet patients' supportive care needs.

Figure A1: Fitch's tiered approach to supportive care



## Appendix B: Psychological needs

Consider a referral to a psychologist, psychiatrist, pastoral/spiritual care practitioner, social worker, specialist nurse or a relevant community-based program if the patient has these issues:

- displaying emotional cues such as tearfulness, distress that requires specialist intervention, avoidance or withdrawal
- being preoccupied with or dwelling on thoughts about cancer and death
- displaying fears about the treatment process or the changed goals of their treatment
- displaying excessive fears about cancer progression or recurrence
- worrying about loss associated with their daily function, dependence on others and loss of dignity
- becoming isolated from family and friends and withdrawing from company and activities that they previously enjoyed
- feeling hopeless and helpless about the effect that cancer is having on their life and the disruption to their life plans
- struggling to communicate with family and loved ones about the implications of their cancer diagnosis and treatment
- experiencing changes in sexual intimacy, libido and function
- struggling with the diagnosis of metastatic or advanced disease
- having difficulties quitting smoking (refer to Quitline on 13 7848) or with other drug and alcohol use
- having difficulties transitioning to palliative care.

Additional considerations that may arise for the multidisciplinary team include:

- support for the carer – encourage referrals to psychosocial support from a social worker, psychologist or general practitioner
- referral to an exercise physiologist or physiotherapist as a therapeutic approach to prevent and manage psychological health
- referral to wellness-after-cancer programs to provide support, information and offer strategies.

# Appendix C: Special population groups

The burden of cancer is not evenly spread across Australia. People experiencing socioeconomic disadvantage, Aboriginal and Torres Strait Islander communities, culturally diverse communities, people living with a disability, people with chronic mental health or psychiatric concerns and those who live in regional and rural areas of Australia have poorer cancer outcomes.

## Aboriginal and Torres Strait Islander people

Cancer is the third leading cause of burden of disease for Aboriginal and Torres Strait Islander people. While Australia's cancer survival rates are among the best in the world, Aboriginal and Torres Strait Islander people continue to experience a different pattern of cancer incidence and significant disparities in cancer outcomes compared with non-Indigenous Australians.

For Aboriginal and Torres Strait Islander people, health and connection to land, culture, community and identity are intrinsically linked. Health encompasses a whole-of-life view and includes a cyclical concept of life–death–life.

The distinct epidemiology of cancer among Aboriginal and Torres Strait Islander people, and unique connection to culture, highlight the need for a specific optimal care pathway for Aboriginal and Torres Strait Islander people with cancer. Ensuring this pathway is culturally safe and supportive is vital to tackling the disparities for Aboriginal and Torres Strait Islander people.

Cancer care for Aboriginal and Torres Strait Islander people should be informed by the Optimal care pathway for Aboriginal and Torres Strait Islander people with cancer, in conjunction with cancer specific optimal care pathways.

In addition to the key principles underpinning cancer-specific pathways, these are the key concepts that are fundamental to Aboriginal and Torres Strait Islander health:

- providing a holistic approach to health and wellbeing
- providing a culturally appropriate and culturally safe service
- acknowledging the diversity of Aboriginal and Torres Strait Islander peoples
- understanding the social determinants and cultural determinants of health (Cancer Australia 2015).

To view the *Optimal care pathway for Aboriginal and Torres Strait Islander people with cancer*, visit the Cancer Australia website <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/OCP-aboriginal-and-torres-strait-islander-people-cancer-second-edition>>. To view the consumer resources – *Checking for cancer* and *Cancer*, visit the Cancer Australia website <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/checking-cancer-what-expect>> and the Cancer Australia website <<https://www.canceraustralia.gov.au/impacted-cancer>>

## People from culturally and linguistically diverse backgrounds

For people from culturally and linguistically diverse backgrounds in Australia, a cancer diagnosis can come with additional complexities, particularly when English proficiency is poor. In many languages there is not a direct translation of the word 'cancer', which can make communicating vital information difficult. Perceptions of cancer and related issues can differ greatly in people from culturally diverse backgrounds, and this can affect their understanding and decision-making after a cancer diagnosis. In addition to different cultural beliefs, when English language is limited, there is potential for miscommunication of important information and advice, which can lead to increased stress and anxiety for patients.

A professionally trained interpreter (not a family member or friend) should be made available when communicating with people with limited English proficiency. Navigation of the Australian health system can pose problems for those with a non-Anglo culture, and members of the treatment teams should pay particular attention to supporting these patients.

The Australian Cancer Survivorship Centre has developed a glossary of more than 700 cancer terms in nine different languages. The multilingual glossary has been designed as a resource for professional translators, interpreters and bilingual health professionals working in the cancer field. The glossary is a unique tool that enables language professionals with access to accurate, consistent and culturally appropriate terminology.

Visit the Peter Mac website <<https://www.petermac.org/about-us/interpreter-services-and-languages/interpreter-service/multilingual-glossary>> to access the glossary.

## People living with disability

Disability, which can be physical, intellectual or psychological, may have existed before the cancer diagnosis or may be new in onset (occurring due to the cancer treatment or incidentally). Adjusting to life with a disability adds another challenge to cancer care and survivorship.

Several barriers prevent people with disabilities from accessing timely and effective health care (AIHW 2017):

- physical limitations
- competing health needs
- the trauma of undergoing invasive procedures
- potential barriers associated with obtaining informed consent
- failure to provide assistance with communication
- lack of information
- discriminatory attitudes among healthcare staff.

In caring for people with disabilities and a cancer diagnosis, the Australian Institute of Health and Welfare disability flag should be used at the point of admittance to correctly identify and meet the additional requirements of a person with disability. Facilities should actively consider access requirements, and health practitioners should make reasonable adjustments where required.

Patients aged between seven and 65 years who have a permanent or significant disability may be eligible for support or funding through the National Disability Insurance Scheme (National Disability Insurance Agency 2018). More information can be found on the NDIS website <[www.ndis.gov.au](http://www.ndis.gov.au)>.

Patients aged 65 years or older (50 years or older for Aboriginal or Torres Strait Islander people) may be eligible for subsidised support and services through aged care services. An application to determine eligibility can be completed online over the phone. More information can be found at the My Aged Care website <[www.myagedcare.gov.au](http://www.myagedcare.gov.au)>.



### More information

'Talking End of Life' is a resource that shows how to teach people with intellectual disability about end of life. It is designed for disability support workers but is also helpful for others including families, health professionals and educators.

To view the resource, visit the Talking End of Life website <[www.caresearch.com.au/tel/tabid/4881/Default.aspx](http://www.caresearch.com.au/tel/tabid/4881/Default.aspx)>.

## Older people with cancer

Planning and delivering appropriate cancer care for older people can present a number of challenges. This could also be true for frail people or those experiencing comorbidities. Effective communication between oncology and geriatrics departments will help facilitate best practice care, which takes into account physiological age, complex comorbidities, risk of adverse events and drug interactions, as well as the implications of cognitive impairment on suitability of treatment and consent (Steer et al. 2009).

At a national interdisciplinary workshop convened by the Clinical Oncology Society of Australia, it was recommended that people over the age of 70 undergo some form of geriatric assessment, in line with international guidelines (COSA 2013; palliAGED 2018). Screening tools can be used to identify those patients in need of a comprehensive geriatric assessment (Decoster et al. 2015). This assessment can be used to help determine life expectancy and treatment tolerance and guide appropriate referral for multidisciplinary intervention that may improve outcomes (Wildiers et al. 2014).

Frailty is not captured through traditional measures of performance status (e.g. ECOG) and includes assessment in the domains of:

- function
- comorbidity
- presence of geriatric syndromes
- nutrition
- polypharmacy
- cognition
- emotional status
- social supports.

## Adolescents and young adults

In recent years, adolescent and young adult oncology has emerged as a distinct field due to lack of progress in survival and quality-of-life outcomes (Ferrari et al. 2010; Smith et al. 2013). The significant developmental change that occurs during this life stage complicates a diagnosis of cancer, often leading to unique physical, social and emotional effects for young people at the time of diagnosis and throughout the cancer journey (Smith et al. 2012).

Cancer care for adolescents and young adults should be informed by the *Optimal care pathway for adolescents and young adults with cancer* <<https://www.cancer.org.au/assets/pdf/optimal-care-pathway-for-adolescents-and-young-adults-with-cancer>>, in conjunction with cancer-specific optimal care pathways.

In caring for young people with cancer, akin to the comorbidities that require specific care in the older cancer population, the treatment team needs to pay careful attention to promoting normal development (COSA 2014). This requires personalised assessments and management involving a multidisciplinary, disease-specific, developmentally targeted approach that adheres to the following principles:

- understanding the developmental stages of adolescence and supporting normal adolescent health and development alongside cancer management
- understanding and supporting the rights of young people
- communication skills and information delivery that are appropriate to the young person
- meeting the needs of all involved, including the young person, their carers and their family
- working with educational institutions and workplaces
- considering survivorship and palliative care needs.

An oncology team caring for an adolescent or young adult with cancer should be able to demonstrate these specific areas of expertise:

- be able to ensure access to expert adolescent and young adult health providers who have knowledge specific to the biomedical and psychosocial needs of the population
- understand the biology and current management of the disease in the adolescent and young adult age group
- consider participating in research and clinical trials for each patient
- engage in proactive discussion and management of fertility preservation, late effects of treatment, ongoing need for contraception, and psychosocial and psychosexual needs
- provide treatment in an environment that is friendly to adolescents and young adults.

## People experiencing socioeconomic disadvantage

In general, people from lower socioeconomic groups are at greater risk of poor health, have higher rates of illness, disability and death, and live shorter lives than those from higher socioeconomic groups (AIHW 2016). People experiencing socioeconomic disadvantage are less likely to participate in screening programs, more likely to be obese, less likely to exercise and much more likely to smoke, which are all risk factors for cancer. In 2012–2016 age-standardised cancer incidence rates were higher in the lowest socioeconomic areas compared with the highest socioeconomic areas for all cancers combined (AIHW 2021).

Socioeconomic status and low health literacy are closely correlated. Therefore, effective communication with patients and carers is particularly important given the prevalence of low health literacy in Australia (estimated at 60 per cent of Australian adults) (ACSQHC 2014).

Consideration should be taken for cancer patients experiencing socioeconomic disadvantage to reduce their risk of being underserved for health care.

## People living with a mental illness

A diagnosis of cancer may present additional challenges to people who have pre-existing chronic mental health or psychiatric concerns, resulting in exacerbation of their mental health symptoms. This may include heightened anxiety, worsening depression or thoughts of self-harm.

As poor adjustment and coping can affect treatment decisions, people who are known to have a mental health diagnosis need psychosocial assessment in the oncology setting to formulate a plan for ongoing support throughout treatment.

Psychosocial support can assist with challenges in communicating with health professionals, enhance understanding of the treatment journey, ensure capacity for consent to treatment options and improve compliance with treatment requests. A referral for psychosocial support from a health professional to the psycho-oncology team can ensure these patients are provided with targeted interventions or referrals to community-based services that may mitigate problems associated with the impacts of social isolation that frequently accompany chronic mental ill-health.

Many patients with chronic mental health problems may be well known to external service providers. Psycho-oncology health professionals can form meaningful partnerships with existing service providers to optimise patient care throughout treatment and beyond.

Drug use disorders fall within the area of mental health conditions. People who are opiate dependent may have specific and individual requirements regarding pain management and their own preference for type of opiate prescribed or used.

## Lesbian, gay, bisexual, transgender, intersex, queer and asexual people

People who identify as sexually or gender diverse may have unique needs following a cancer diagnosis. Sexually or gender diverse identities include (but are not limited to) people who identify as lesbian, gay, bisexual, transgender, intersex, queer and asexual collectively 'LGBTIQ+'.

Sexual orientation and gender identity are relevant across the entire spectrum of cancer care, from prevention to survivorship and end-of-life care. LGBTIQ+ people are less likely to participate in cancer screening, and some segments of the LGBTIQ+ community exhibit elevated rates of specific cancer risk factors – for example, higher rates of smoking and alcohol use. Regarding treatment, there may be unique factors relevant to LGBTIQ+ people that may affect decision-making. Additionally, the LGBTIQ+ population experiences higher rates of anxiety, depression and stressful life circumstances, and may be at risk of inferior psychosocial outcomes following a cancer diagnosis. LGBTIQ+ people are also more likely to be estranged from their families of origin, and for older people, less likely to have adult children who may provide support and care.

Barriers to care for LGBTIQ+ people include past negative interactions with health systems, experiences or fear of discrimination and harassment in healthcare settings, assumptions of cisgender/heterosexual identity, lack of recognition or exclusion of same-sex partners from care, and a lack of relevant supportive care and information resources.

To provide safe and appropriate care for LGBTIQ+ people with cancer, healthcare providers should:

- display environmental cues to show an inclusive and safe setting for LGBTIQ+ patients
- avoid assumptions about the sexual orientation or gender identity of patients and their partners
- facilitate positive disclosure of sexual orientation or gender identity
- include same-sex/gender partners and families of choice in care
- be aware of relevant supportive care and information resources
- provide non-judgemental, patient-centred care.

## People who live in regional, rural and remote areas

People living in remote areas of Australia are 1.3 times more likely to die from cancer and have a lower five-year relative survival rate compared with those living in major cities (AIHW 2021). Also, people living in rural and remote areas have higher rates of hospitalisation, death and injury and have poorer access to and use of primary health care services than people living in major cities.

Poorer cancer outcomes may be related to health workforce shortages in remote and very remote areas, despite having a greater need for medical services and practitioners with a broader scope of practice.

People living in remote areas are also more likely to engage in risky behaviours, such as smoking and consuming alcohol at excessive levels, and are less likely to take part in cancer screening services compared with those living in major cities.

People living in rural and remote areas face barriers to accessing health care due to challenges of geographic spread, low population density, limited infrastructure and the higher costs of delivering rural and remote health care. Travel is a significant barrier to accessing health services and can lead to reduced access to cancer care across the care pathway. Other costs associated with travel to metropolitan centres increases the impact of a cancer diagnosis on a person and their families.

To improve the cancer experience and outcomes for people who live in rural and remote areas of Australia the entire health system must address:

- improved access to preventive care in rural and remote areas, including more accessible and affordable options through pop-up services and virtual care
- extra support through increased access to healthy eating choices that address issues of food security, and funded programs to support education and lifestyle behaviour change
- tailored education and awareness campaigns to increase health literacy
- improved access to health care, income, housing and food security on cancer outcomes for people living in rural and remote areas
- risk-based population cancer screening to include the social and geographical risks that increase the likelihood of developing cancer for people living in rural and remote locations, including reduced access to regular health check-ups and screening, sun exposure, food insecurity and higher rates of smoking
- assessment of risk-based population screening initiatives to consider scalability into rural and remote areas to ensure equitable access (Cancer Australia 2023).

## Appendix D: Complementary therapies

Complementary therapies may be used together with conventional medical treatments to support and enhance quality of life and wellbeing. They do not aim to cure the patient's cancer. Instead, they are used to help control symptoms such as pain and fatigue (Cancer Council Australia 2019).

The lead clinician or health professional involved in the patient's care should discuss the patient's use (or intended use) of complementary therapies not prescribed by the multidisciplinary team to assess safety and efficacy and to identify any potential toxicity or drug interactions.

The lead clinician should seek a comprehensive list of all complementary and alternative medicines being taken and explore the patient's reason for using these therapies and the evidence base. A transparent and honest discussion that is free from judgement should be encouraged.

While some complementary therapies are supported by strong evidence, others are not. For such therapies, the lead clinician should discuss their potential benefits and use them alongside conventional therapies (NHMRC 2014).

If the patient expresses an interest in using complementary therapies, the lead clinician should consider referring patients to health providers within the multidisciplinary team who have expertise in the field of complementary and alternative therapies (e.g. a clinical pharmacist, dietitian or psychologist) to assist them to reach an informed decision. Costs of such approaches should be part of the discussion with the patient and considered in the context of evidence of benefit.

The lead clinician should assure patients who use complementary therapies that they can still access a multidisciplinary team review and encourage full disclosure about therapies being used.



### More information

Refer to the Clinical Oncological Society of Australia's position statement Use of complementary and alternative medicine by cancer patients <[https://www.cosa.org.au/media/1133/cosa\\_cam-position-statement\\_final\\_new-logo.pdf](https://www.cosa.org.au/media/1133/cosa_cam-position-statement_final_new-logo.pdf)>.

# Appendix E: Members of the multidisciplinary team for biliary cancer

The multidisciplinary team may include the following members:

- dietitian\*
- hepatopancreatobiliary surgeon\*
- interventional endoscopist (gastroenterologist or surgeon)\*
- medical oncologist\*
- pathologist\*
- radiation oncologist\*
- upper GI cancer nurse consultant/coordinator\*
- upper GI radiologist (diagnostic and interventional)\*
- hepatologist
- Aboriginal health practitioner, Indigenous liaison officer or remote general practitioner
- anaesthetist
- exercise physiologist
- fertility specialist
- general practitioner
- geneticist
- nuclear medicine physician
- physiotherapist
- palliative care
- social worker
- psychologist
- spiritual/pastoral care.

\* Denotes core members. Core members of the multidisciplinary team are expected to attend most multidisciplinary team meetings either in person or remotely.

# Resource list

## For patients, families and carers

### Advance Care Planning Australia

Advance Care Planning Australia provides national advance care planning resources for individuals, families, health professional and service providers. Resources include a national advisory service, information for each state and territory, and online education for health professionals.

- Telephone: 1300 208 582
- Website <[www.advancecareplanning.org.au](http://www.advancecareplanning.org.au)>

### Australian Cancer Survivorship Centre

The Australian Cancer Survivorship Centre has developed information resources and events to help people move from initial treatment to post treatment and beyond, including those receiving maintenance treatments. While they do not provide clinical advice, they connect with a range of providers to enable improved care.

- Telephone: (03) 8559 6220
- Website <[www.petermac.org/cancersurvivorship](http://www.petermac.org/cancersurvivorship)>

### Australian Commission on Safety and Quality in Health Care

The Australian Commission on Safety and Quality in Health Care has developed a resource for patients and carers explaining the coordination of care that patients should receive from their health service during cancer treatment. The resource is called *What to expect when receiving medication for cancer care* <<https://www.safetyandquality.gov.au/publications-and-resources/resource-library/what-expect-when-receiving-medication-cancer-care-fact-sheet>>.

### Beyond Blue

Beyond Blue provides information about depression, anxiety and related disorders, as well as about available treatment and support services.

- Telephone: 1300 22 4636
- Website <[www.beyondblue.org.au](http://www.beyondblue.org.au)>

### Cancer Council's Cancer Information and Support Service

Cancer Council 13 11 20 is a confidential telephone support service available to anyone affected by cancer. This service acts as a gateway to evidence-based documented, practical and emotional support available through Cancer Council services and other community organisations. Calls will be answered by a nurse or other oncology professional who can provide information relevant to a patient's or carer's situation. Health professionals can also access this service.

- Telephone: 13 11 20 – Monday to Friday, 9 am to 5 pm (some states have extended hours)
- Website <[www.cancer.org.au/about-us/state-and-territory-councils/](http://www.cancer.org.au/about-us/state-and-territory-councils/)>

### Cancer Council's Cancer Connect

Cancer Connect is a free and confidential telephone peer support service that connects someone who has cancer with a specially trained volunteer who has had a similar cancer experience.

A Connect volunteer can listen with understanding and share their experiences and ways of coping. They can provide practical information, emotional support and hope. Many people newly diagnosed with cancer find this one-to-one support very beneficial.

For more information on Cancer Connect call Cancer Council on 13 11 20.

## Canteen

Canteen helps adolescents, young adults and parents to cope with cancer in their family. Canteen offers individual support services, peer support services and a youth cancer service, as well as books, resources and useful links.

- Telephone: 1800 835 932 to talk to a health professional about information and support for young people or 1800 226 833 for other enquiries
- Website <[www.canteen.org.au/](http://www.canteen.org.au/)>

## Carer Gateway

Carer Gateway is an Australian Government program providing free services and support for carers. Carer Gateway service providers help carers access a range of free services and support just for carers. These services can be accessed in-person, over the phone or online through the website.

- Telephone: 1800 422 737 to find out if you are eligible to access services offered by the team of Carer Gateway service providers across Australia.
- Website <[www.carergateway.gov.au](http://www.carergateway.gov.au/)>

## Clinical trial information

For a collection of clinical trials available in Australia refer to the following sources of information:

- Australian clinical trials <[www.australianclinicaltrials.gov.au](http://www.australianclinicaltrials.gov.au/)>
- Australian New Zealand Clinical Trials Registry <[www.anzctr.org.au](http://www.anzctr.org.au/)>
- ClinicalTrials.gov <[www.clinicaltrials.gov](http://www.clinicaltrials.gov/)>.

## CanEAT pathway

A guide to optimal cancer nutrition for people with cancer, carers and health professionals.

- Webpage on malnutrition <[https://education.eviq.org.au/malnutrition-sarcopenia](https://education.eviq.org.au/malnutrition-sarcopenia/)>
- Webpage on eating well <[https://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/eating-well](https://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/eating-well/)>

## Guides to best cancer care

The short guides help patients, carers and families understand the optimal cancer care that should be provided at each step. They include optimal timeframes within which tests or procedures should be completed, prompt lists to support patients to understand what might happen at each step of their cancer journey and to consider what questions to ask, and provide information to help patients and carers communicate with health professionals.

The guides are located on an interactive web portal, with downloadable PDFs available in multiple languages.

- Website <[https://www.cancer.org.au/cancercareguides](https://www.cancer.org.au/cancercareguides/)>

## Liver Foundation

The Liver Foundation's free nurse-led support line (1800 841 118) provides expert guidance for people with biliary cancer, primary liver cancer and underlying liver conditions. The foundation offers reliable, easy-to-understand information about diagnosis, treatment and managing your health. You can also access practical resources on the website.

- Telephone: 1800 841 118
- Website <[https://liver.org.au](https://liver.org.au/)>

## Look Good, Feel Better

A free national community service program, run by the Cancer Patients Foundation, dedicated to teaching cancer patients how to manage the appearance-related side effects caused by treatment for any type of cancer.

- Telephone: 1800 650 960
- Website <[https://lgfb.org.au](https://lgfb.org.au/)>

## Pancare Foundation

Pancare Foundation is a leading charity committed to inspiring hope, raising awareness, supporting families and funding research for upper gastrointestinal (GI) cancers. With some of the lowest survival rates out of all the major cancers, Pancare is dedicated to improving outcomes for those affected by upper GI cancers including pancreatic, liver, stomach, biliary and oesophageal cancers.

Pancare provides support, resources and information for people living with cancer and their families from the point of diagnosis through to treatment and beyond.

- Telephone: 1300 881 698
- Website <<https://pancare.org.au>>

## Quitline

Quitline is a confidential, evidence-based telephone counselling service. Highly trained Quitline counsellors use behaviour change techniques and motivational interviewing over multiple calls to help people plan, make and sustain a quit attempt.

Quitline is a culturally inclusive service for all, and Aboriginal counsellors are also available. Health professionals can refer patients to Quitline online or via fax.

- Telephone: 13 7848
- Website <[www.quit.org.au](http://www.quit.org.au)> or the relevant website in your state or territory.

## For health providers

### Australian Cancer Survivorship Centre

The Australian Cancer Survivorship Centre provides expertise in survivorship care, information, support and education. Its purpose is to support and enable optimal survivorship care.

- Telephone: (03) 8559 6220
- Website <[www.petermac.org/cancersurvivorship](http://www.petermac.org/cancersurvivorship)>

## Australian Commission on Safety and Quality in Health Care

The Australian Commission on Safety and Quality in Health Care has developed a guide for clinicians containing evidence-based strategies to support clinicians to understand and fulfil their responsibilities to cancer patients. This guide is particularly relevant to steps 3 to 6 of the optimal care pathway. The guide is titled *NSQHS Standards user guide for medication management in cancer care for clinicians* <<https://www.safetyandquality.gov.au/publications-and-resources/resource-library/nsqhs-standards-user-guide-medication-management-cancer-care-clinicians>>.

## Cancer Council Australia

Information on prevention, research, treatment and support provided by Australia's peak independent cancer authority.

- Website <[www.cancer.org.au](http://www.cancer.org.au)>

## CanEAT pathway

A guide to optimal cancer nutrition for people with cancer, carers and health professionals.

- Webpage on malnutrition <<https://education.eviq.org.au/malnutrition-sarcopenia>>
- Webpage on eating well <<https://patients.cancer.nsw.gov.au/coping-with-cancer/physical-wellbeing/eating-well>>.

## eviQ

A clinical information resource providing health professionals with current evidence-based, peer-maintained, best practice cancer treatment protocols and information relevant to the Australian clinical environment.

- Website <[www.eviq.org.au](http://www.eviq.org.au)>

## **National Health and Medical Research Council**

Information on clinical practice guidelines, cancer prevention and treatment.

- Website <[www.nhmrc.gov.au](http://www.nhmrc.gov.au)>

## **Pam McLean Centre**

The Pam McLean Centre offers workshops for health professionals to develop their ability to communicate well with different stakeholders, build good communication skills and habits, and increase their confidence in creating an environment for good communication between all parties.

- Website <<https://pammcleancentre.org/>>

# Glossary

**advance care directive** – voluntary person-led document that focus on an individual's values and preferences for future health and medical treatment decisions, preferred outcomes and care. They are completed and signed by a competent person. They are recognised by specific legislation (statutory) or common law (non-statutory). Each state and territory has different terminology and legislation surrounding advance care directives and substitute decision-makers. A person can appoint the substitute decision-maker(s) who can make decisions about health or personal care on the individual's behalf if they are no longer able to make decisions themselves. In some states and territories this appointment is within the advance care directive form; in others it requires a separate form. Advance care directives focus on the future health care of a person, not on the management of his or her assets. They come into effect when an individual loses decision-making capacity.

**advance care planning** – the process of planning for future health and personal care, where the person's values, beliefs and preferences are made known so they can guide decision-making at a future time when that person cannot make or communicate their decisions.

**alternative therapies** – treatments used in place of conventional medical treatment.

**care coordinator** – the health provider nominated by the multidisciplinary team to coordinate patient care. The care coordinator may change over time depending on the patient's stage in the care pathway and the location and care in which care is being delivered.

**complementary therapies** – supportive treatment used in conjunction with conventional medical treatment. These treatments may improve wellbeing and quality of life and help people deal with the side effects of cancer.

**end-of-life care** – includes physical, spiritual and psychosocial assessment, and care and treatment, delivered by health professionals and ancillary staff. It also includes support of families and carers and care of the patient's body after their death.

**immunotherapy** – a type of cancer treatment that helps the body's immune system to fight cancer. Immunotherapy can boost the immune system to work better against cancer or remove barriers to the immune system attacking the cancer.

**indicator** – a documentable or measurable piece of information regarding a recommendation in the optimal care pathway.

**informed financial consent** – the provision of cost information to patients, including notification of likely out-of-pocket expenses (gaps), by all relevant service providers, preferably in writing, before admission to hospital or treatment (Commonwealth Department of Health 2017).

**lead clinician** – the clinician who is nominated as being responsible for individual patient care. The lead clinician may change over time depending on the stage of the care pathway and where care is being provided.

**metastatic disease** – cancer that has spread from the part of the body where it started (the primary site) to other parts of the body.

**multidisciplinary care** – an integrated team approach to health care in which medical and allied health providers consider all relevant treatment options and collaboratively develop an individual treatment plan for each patient.

**multidisciplinary team** – comprises the core disciplines that are integral to providing good care. The team is flexible in approach, reflects the patient's clinical and psychosocial needs and has processes to facilitate good communication.

**multidisciplinary team meeting** – a meeting of health professionals from one or more clinical disciplines who together make decisions about recommended treatment of patients.

**optimal care pathway** – the key principles and practices required at each stage of the care pathway to guide the delivery of consistent, safe, high-quality and evidence-based care for all people affected by cancer.

**performance status** – an objective measure of how well a patient can carry out activities of daily life.

**primary care health professional** – in most cases this is a general practitioner but may also include general practice nurses, community nurses, nurse practitioners, allied health professionals, midwives, pharmacists, dentists and Aboriginal health workers.

**spiritual care** – the aspect of humanity that refers to the way individuals seek and express meaning and purpose and the way they experience their connectedness to the moment, to self, to others, to nature, and to the significant or sacred.

**substitute decision-maker** – a person permitted under the law to make decisions on behalf of someone who does not have competence or capacity.

**supportive care** – care and support that aims to improve the quality of life of people living with cancer, cancer survivors and their family and carers and particular forms of care that supplement clinical treatment modalities.

**survivorship** – an individual is considered a cancer survivor from the time of diagnosis, and throughout their life; the term includes individuals receiving initial or maintenance treatment, in recovery or in the post-treatment phase.

**survivorship care plan** – a formal, written document that provides details of a person's cancer diagnosis and treatment, potential late and long-term effects arising from the cancer and its treatment, recommended follow-up, surveillance, and strategies to remain well.

**targeted therapy** – a medicine that blocks the growth and spread of cancer by interfering with specific molecules.

**voluntary assisted dying** – the assistance provided to a person by a health practitioner to end their life. 'Voluntary' indicates that the practice is a voluntary choice of the person and that they are competent (have capacity) to decide to access voluntary assisted dying.

## References

- Aloia TA, Járufe N, Javle M, Maithel SK, Roa JC, Adsay V, et al. 2015, 'Gallbladder cancer: expert consensus statement', *HPB* (Oxford), vol. 17, no. 8, pp. 681–690.
- Australian Adult Cancer Pain Management Guideline Working Party 2019, *Australian adult cancer pain management guideline: 'Cancer pain management in adults'*, Cancer Council Australia, Sydney, viewed 17th June 2025, <<https://www.cancer.org.au/assets/pdf/archived-cancer-pain-management-in-adults-guidelines-19-july-2019>>.
- Australian Cancer Survivorship Centre 2024, '*Survivorship care planning*', viewed 17th June 2025, <<https://www.petermac.org/document/survivorship-care-planning-initiative/viewdocument/1097>>.
- Australian Cancer Survivorship Centre 2021, *Community support organisations' cancer survivorship care consensus statement version 3*, viewed 17th June 2025, <<https://www.petermac.org/component/edocman/ngo-consensusstatement-v3-sept-2021-web/viewdocument/516?Itemid=0>>.
- Australian Clinical Trials 2015, '*Potential benefits and risks*', National Health and Medical Research Council, Department of Industry, Innovation and Science, Australian Government, Canberra, viewed 17 June 2025, <<https://www.australianclinicaltrials.gov.au/participants/what-you-should-know#potential-benefits-and-risks>>.
- Australian Commission on Safety and Quality in Health Care (ACSQHC) 2014, *Health literacy: taking action to improve safety and quality*, ACSQHC, Sydney, viewed 18 February 2020, <<https://www.safetyandquality.gov.au/sites/default/files/migrated/Health-Literacy-Taking-action-to-improve-safety-and-quality.pdf>>.
- Australian Commission for Safety and Quality in Health Care (ACSQHC) 2015, *Credentialing health practitioners and defining their scope of clinical practice: a guide for managers and practitioners*, ACSQHC, Sydney, viewed 18 February 2020, <<https://www.safetyandquality.gov.au/publications-and-resources/resource-library/credentialing-health-practitioners-and-defining-their-scope-clinical-practice-guide-managers-and-practitioners>>.
- Australian Commission on Safety and Quality in Health Care (ACSQHC) 2017, *National Safety and Quality Health Service Standards guide for hospitals*, ACSQHC, Sydney, viewed 18 February 2020, <<https://www.safetyandquality.gov.au/wp-content/uploads/2017/12/National-Safety-and-Quality-Health-Service-Standards-Guide-for-Hospitals.pdf>>.
- Australian Commission on Safety and Quality in Health Care (ACSQHC) 2019a, *Person-centred care*, ACSQHC, Sydney, viewed 15 May 2020, <<https://www.safetyandquality.gov.au/our-work/partnering-consumers/person-centred-care>>.
- Australian Commission on Safety and Quality in Health Care (ACSQHC) 2019b, *Australian Hospital Patient Experience Question Set*, ACSQHC, Sydney, viewed 25 March 2020, <<https://www.safetyandquality.gov.au/our-work/indicators-measurement-and-reporting/australian-hospital-patient-experience-question-set>>.
- Australian Commission for Safety and Quality in Health Care (ACSQHC) 2020, '*NSQHS Standards user guide for medication management in cancer care*', ACSQHC, Sydney, viewed 16 April 2020, <<https://www.safetyandquality.gov.au/publications-and-resources/resource-library/nsqhs-standards-user-guide-medication-management-cancer-care>>.
- Australian Institute of Health and Welfare (AIHW) 2016, *Australia's health 2016*, Australia's health series no. 15. Cat. no. AUS 199, AIHW, Canberra.

Australian Institute of Health and Welfare (AIHW) 2017, *Access to health services by Australians with disability*, viewed 17 June 2025, <<https://www.aihw.gov.au/reports/disability/access-health-services-disability/contents/summary>>.

Australian Institute of Health and Welfare (AIHW) 2018, *Australia's health 2018*, Australia's health series no. 16. Cat. no. AUS 221, AIHW, Canberra.

Australian Institute of Health and Welfare (AIHW) 2021, *Cancer data in Australia 2021*, Cancer series no.133. Cat. no. CAN 144, AIHW, Canberra.

AYA Cancer Fertility Preservation Guidance Working Group 2014, *Fertility preservation for AYAs diagnosed with cancer: guidance for health professionals*, Cancer Council Australia, Sydney, viewed 20 July 2020, <[https://wiki.cancer.org.au/australia/COSA:AYA\\_cancer\\_fertility\\_preservation/Introduction](https://wiki.cancer.org.au/australia/COSA:AYA_cancer_fertility_preservation/Introduction)>.

Baria K, De Toni EN, Yu B, Jiang Z, Kabadi SM, Malvezzi M 2022, 'Worldwide incidence and mortality of biliary tract cancer', *Gastro Hep Advances*, vol. 1, no. 4, pp. 618–626.

Bridgewater J, Galle P, Khan S, Patel T, Pawlik T, Gores G 2014. 'Guidelines for the diagnosis and management of intrahepatic cholangiocarcinoma', *Journal of Hepatology*, vol. 60, no. 6, pp. 1268–1289.

Cancer Australia 2015, *National Aboriginal and Torres Strait Islander cancer framework*, Cancer Australia, Surry Hills, viewed 24 August 2017, <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/national-aboriginal-and-torres-strait-islander-cancer-framework>>.

Cancer Australia 2017, *Principles of cancer survivorship*, Australian Government, Sydney, viewed 17 June 2025, <[https://www.canceraustralia.gov.au/sites/default/files/migrated-files/publications/principles-cancer-survivorship/pdf/pocs\\_-\\_principles\\_of\\_cancer\\_survivorship.pdf](https://www.canceraustralia.gov.au/sites/default/files/migrated-files/publications/principles-cancer-survivorship/pdf/pocs_-_principles_of_cancer_survivorship.pdf)>.

Cancer Australia 2019, *Principles of multidisciplinary care*, Australian Government, Sydney, viewed 17 June 2025, <<https://www.canceraustralia.gov.au/clinicians-hub/multidisciplinary-care/all-about-multidisciplinary-care/principles-multidisciplinary-care>>.

Cancer Australia 2020, *Cancer care in the time of COVID-19: a conceptual framework for the management of cancer during a pandemic*, Australian Government, Sydney, viewed 3 September 2020, <<https://www.canceraustralia.gov.au/publications-and-resources/cancer-australia-publications/cancer-care-time-covid-19-conceptual-framework-management-cancer-during-pandemic>>.

Cancer Australia 2023, *Australian Cancer Plan*, Australian Government, Sydney, viewed 12 December 2024, <<https://www.australiacancerplan.gov.au/populations/people-living-in-rural-and-remote-areas>>.

Cancer Council Australia 2024, '*Prevention*', Cancer Council Australia, Sydney, viewed 11 December 2024, <<https://www.cancer.org.au/cancer-information/causes-and-prevention>>.

Cancer Council Australia 2020, '*Facing end of life*', Cancer Council Australia, Sydney, viewed 3 May 2025, <<https://www.cancer.org.au/assets/pdf/facing-end-of-life>>.

Cancer Council Australia 2019, '*Complementary and alternative therapies*', Cancer Council Australia, Sydney, viewed 22 July 2019, <<https://www.cancer.org.au/about-cancer/treatment/complementary-therapies-and-cancer.html>>.

Cancer Council Victoria 2019, '*Palliative care*', Cancer Council Victoria, viewed 7 October 2019, <[https://www.cancervic.org.au/cancer-information/treatments/treatments-types/palliative\\_care/palliative-care-treatment.html](https://www.cancervic.org.au/cancer-information/treatments/treatments-types/palliative_care/palliative-care-treatment.html)>.

- Cancer Research in Primary Care 2016, *Principles statement: Shared care*, PC4 Shared Care Working Group, Melbourne, viewed 4 October 2019, <<http://pc4tg.com.au/wp-content/uploads/2016/07/PC4-Principles-Statement-shared-care-2016-1.pdf>>.
- Clinical Oncology Society of Australia (COSA) 2013, Special Issue: COSA's 40th Annual Scientific Meeting, Cancer Care Coming of Age, 12–14 November 2013, Adelaide Convention Centre, *Asia-Pacific Journal of Clinical Oncology*, vol. 9, no. 3, pp. 61–98.
- Clinical Oncology Society of Australia (COSA) 2014, *Psychosocial management of AYAs diagnosed with cancer: guidance for health professionals*, COSA, Sydney, viewed 7 October 2019, <[http://wiki.cancer.org.au/australia/COSA:Psychosocial\\_management\\_of\\_AYA\\_cancer\\_patients](http://wiki.cancer.org.au/australia/COSA:Psychosocial_management_of_AYA_cancer_patients)>.
- Clinical Oncology Society of Australia (COSA) 2015, *Cancer care coordinator: position statement*, COSA, Sydney, viewed 22 July 2019, <[https://www.cosa.org.au/media/332296/cancer-care-coordinator-position-statement\\_final-endorsed-by-council\\_161115\\_cnsa-logo.pdf](https://www.cosa.org.au/media/332296/cancer-care-coordinator-position-statement_final-endorsed-by-council_161115_cnsa-logo.pdf)>.
- Clinical Oncology Society of Australia (COSA) 2016, *Australasian tele-trial model: access to clinical trials closer to home using tele-health a national guide for implementation*, v1.7, COSA, Sydney, viewed 18 July 2019, <<https://www.cosa.org.au/media/332325/cosa-teletrial-model-final-19sep16.pdf>>.
- Clinical Oncology Society of Australia (COSA) 2018, *COSA position statement on exercise in cancer care*, COSA, Sydney, viewed 22 July 2019, <<https://www.cosa.org.au/media/332488/cosa-position-statement-v4-web-final.pdf>>.
- Commonwealth Department of Health 2017, *Out-of-pocket expenses for private medical treatment (informed financial consent)*, Commonwealth of Australia, Canberra.
- Cormie P, Atkinson M, Bucci L, Cust A, Eakin E, Hayes S, et al. 2018, 'Clinical Oncology Society of Australia position statement on exercise in cancer care', *Medical Journal of Australia*, vol. 209, no. 4, pp. 184–187.
- Cormie P, Zopf EM, Zhang X, Schmitz KH 2017, 'The impact of exercise on cancer mortality, recurrence, and treatment-related adverse effects', *Epidemiologic Reviews*, vol. 39, no. 1, pp. 71–92.
- Decoster L, Van Puyvelde K, Mohile S, Wedding U, Basso U, Colloca G, et al. 2015, 'Screening tools for multidimensional health problems warranting a geriatric assessment in older cancer patients: an update on SIOG recommendations', *Annals of Oncology*, vol. 26, no.1, pp 288–300.
- European Association for the Study of the Liver 2016, 'Clinical practice guidelines on the prevention, diagnosis and treatment of gallstones', *Journal of Hepatology*, vol. 65, no. 1, pp. 146–181.
- European Association for the Study of the Liver 2023, 'EASL-ILCA clinical practice guidelines on the management of intrahepatic cholangiocarcinoma', *Journal of Hepatology*, vol. 79, no. 1, pp. 181–208.
- Emery J 2014, 'Cancer survivorship – the role of the GP', *Australian Family Physician*, vol. 43, no. 8, pp. 521–525.
- eviQ 2022, 'Safe handling and waste management of hazardous drugs', Cancer Institute NSW, Sydney, viewed 11 December 2024, <<https://www.eviq.org.au/clinical-resources/administration-of-antineoplastic-drugs/188-safe-handling-and-waste-management-of-hazardou>>.
- Ferrari A, Thomas D, Franklin A, Hayes-Lattin B, Mascarin M, van der Graaf W, et al. 2010, 'Starting an adolescent and young adult program: some success stories and some obstacles to overcome', *Journal of Clinical Oncology*, vol. 28, no. 32, pp. 4850–4857.

- Fitch M 2000, 'Supportive care for cancer patients', *Hospital Quarterly*, vol. 3, no. 4, pp. 39–46.
- Fitch MI 2008, 'Supportive care framework', *Canadian Oncology Nursing Journal*, vol. 18, no. 1, pp. 6–14.
- GI Cancer Institute 2024, 'Gallbladder & biliary tract cancer', viewed 11 December 2024, <<https://gicancer.org.au/cancer/gallbladder-and-biliary-cancer/>>.
- Gilligan T, Coyle N, Frankel RM, Berry DL, Bohlke K, Epstein, RM, et al. 2017, 'Patient–clinician communication: American Society of Clinical Oncology consensus guideline', *Journal of Clinical Oncology*, vol. 35, no. 31, pp. 3618–3623.
- Hack TF, Reuther DJ, Weir LM, Grenier D, Degner LF 2012, 'Promoting consultation recording practice in oncology: identification of critical implementation factors and determination of patient benefit', *Psycho-Oncology*, vol. 22, pp. 1273–1282.
- Haines IE 2011, 'Managing patients with advanced cancer: the benefits of early referral for palliative care', *Medical Journal of Australia*, vol. 194, no. 3, pp. 107–108.
- Hayes SC, Newton RU, Spence RR, Galvao DA 2019, 'The Exercise and Sports Science Australia position statement: exercise medicine in cancer management', *Journal of Science and Medicine in Sport*, vol. 22, no. 11, pp. 1175–1199.
- Hewitt M, Greenfield S, Stovall E 2006, *From cancer patient to cancer survivor: lost in transition*, National Academies Press, Washington.
- Jefford M, Ward AC, Lisy K, Lacey K, Emery JD, Glaser AW, et al. 2017, 'Patient-reported outcomes in cancer survivors: a population-wide cross-sectional study', *Support Care Cancer*, no. 10, pp. 3171–3179.
- Khan SA, Tavolari S, Brandi G 2019, 'Cholangiocarcinoma: epidemiology and risk factors', *Liver International*, no. 39 (Suppl. 1), pp. 19–31.
- Kim HS, Kang MJ, Kang J, Kim K, Kim B, Kim SH, et al. 2024, 'Practice guidelines for managing extrahepatic biliary tract cancers', *Annals of Hepato-Biliary-Pancreatic Surgery*, vol. 28, no. 2, pp. 161–202.
- Laidsaar-Powell R, Butow P, Boyle F, Juraskova 2018a, 'Facilitating collaborative and effective family involvement in the cancer setting: guidelines for clinicians (TRIO guidelines-1)', *Patient Education and Counseling*, vol. 101, no. 6, pp. 970–982.
- Laidsaar-Powell R, Butow P, Boyle F, Juraskova 2018b, 'Managing challenging interactions with family caregivers in the cancer setting: guidelines for clinicians (TRIO guidelines-2)', *Patient Education and Counselling*, vol. 101, no. 6, pp. 983–994.
- Lisy K, Langdon L, Piper A, Jefford M 2019, 'Identifying the most prevalent unmet needs of cancer survivors in Australia: a systematic review', *Asia-Pacific Journal of Clinical Oncology*, vol. 15, no. 5, pp. e68–e78.
- Makki M, Bentaleb M, Abdulrahman M, Suhood A, Al Harthi S, Ribeiro A Jr, 2024, 'Current interventional options for palliative care for patients with advanced-stage cholangiocarcinoma', *World Journal of Clinical Oncology*, vol. 15, no. 3, pp. 381–390.
- Mansour JC, Aloia TA, Crane CH, Heimbach JK, Nagino M, Vauthey JN 2015. 'Hilar cholangiocarcinoma: expert consensus statement', *HPB (Oxford)*, vol. 17, no. 8, pp. 691–699.
- Monterosso L, Platt V, Bulsara M, Berg M 2019, 'Systematic review and meta-analysis of patient reported outcomes for nurse-led models of survivorship care for adult cancer patients', *Cancer Treatment Reviews Journal*, no. 73, pp. 62–72.

- My Health Record 2019, 'What is My Health Record?' Australian Government, Sydney, viewed 17 December 2019, <<https://www.myhealthrecord.gov.au/for-you-your-family/what-is-my-health-record>>.
- Nagino M, Hirano S, Yoshitomi H, Aoki T, Uesaka K, Unno M, et al. 2021, 'Clinical practice guidelines for the management of biliary tract cancers', 3rd English edition, *Journal of the Hepatobiliary Pancreatic Sciences*, vol. 28, no. 1, pp. 26–54.
- National Disability Insurance Agency 2018, 'How the NDIS works', NDIS, Canberra, viewed 3 June 2019, <<https://www.ndis.gov.au/understanding/how-ndis-works>>.
- National Health and Medical Research Council (NHMRC) 2013, *Personalised medicine and genetics*, viewed 3 June 2019 <<https://www.nhmrc.gov.au/about-us/publications/personalised-medicine-and-genetics>>.
- National Health and Medical Research Council (NHMRC) 2014, *Talking with your patients about complementary medicine: a resource for clinicians*, NHMRC, Canberra, viewed 18 February 2020, <<https://www.caresearch.com.au/caresearch/ClinicalPractice/PatientConsiderations/ComplementaryTherapies/tabid/1258/Default.aspx>>.
- Palepu J, Endo I, Chaudhari V, Murthy GVS, Chaudhuri S, Adam R, et al. 2025, 'IHPBA-APHPBA clinical practice guidelines': international Delphi consensus recommendations for gallbladder cancer, *HPB*, vol. 26, no. 11, pp. 1311–1326.
- palliAGED 2018, 'Needs Assessment', Flinders University, Bedford Park, viewed 1 October 2019, <<https://www.palliaged.com.au/tabid/4879/Default.aspx>>.
- Palliative Care Australia 2024, *National Palliative Care Standards*, 5.1 edn, Palliative Care Australia, Canberra, viewed 11 December 2024, <<https://palliativecare.org.au/national-palliative-care-standards-for-specialist-palliative-care-providers-5-1-edition-2024/>>.
- Palliative Care Victoria 2025, 'Spiritual care', Palliative Care Victoria, Melbourne, viewed 17 June 2025, <<https://www.pallcarevic.asn.au/page/143/spiritual-care>>.
- Queensland Health 2015, *Sad news sorry business: guidelines for caring for Aboriginal and Torres Strait Islander people through death and dying*, State Government of Queensland, Brisbane, viewed 22 July 2019, <[https://www.health.qld.gov.au/\\_\\_data/assets/pdf\\_file/0023/151736/sorry\\_business.pdf](https://www.health.qld.gov.au/__data/assets/pdf_file/0023/151736/sorry_business.pdf)>.
- Ramjeesingh R, Chaudhury P, Tam VC, Roberge D, Lim HJ, Knox JJ, et al. 2023, 'A practical guide for the systemic treatment of biliary tract cancer in Canada', *Current Oncology*, vol. 30, no. 8, pp. 7132–7150.
- Roth GS, Verlingue L, Sarabi M, Blanc JF, Boleslawski E, Boudjema K, et al. 2024, 'Biliary tract cancers: French national clinical practice guidelines for diagnosis, treatments and follow-up', *European Journal of Cancer*, no. 202, p. 114000.
- Shroff RT, Kennedy EB, Bachini M, Bekaii-Saab T, Crane C, Edeline J, et al. 2019, 'Adjuvant therapy for resected biliary tract cancer: ASCO Clinical Practice Guideline', *Journal of Clinical Oncology*, vol. 37, no. 12, pp. 1015–1027.
- Silver JK, Baima J 2013, 'Cancer prehabilitation: an opportunity to decrease treatment-related morbidity, increase cancer treatment options, and improve physical and psychological health outcomes', *American Journal of Physical Medicine & Rehabilitation*, vol. 92, no. 8, pp. 715–727.

- Silver JK 2015, 'Cancer prehabilitation and its role in improving health outcomes and reducing health care costs', *Seminars in Oncology Nursing*, vol. 31, no. 1, pp. 1–3.
- Sjoquist K, Zalcberg J 2013, 'Clinical trials – advancing cancer care', *Cancer Forum*, vol. 37, no. 1, pp. 80–88.
- Smith A, Bellizzi K, Keegan T, Zebrack B, Chen V, Neale A, et al. 2013, 'Health-related quality of life of adolescent and young adult patients with cancer in the United States: the Adolescent and Young Adult Health Outcomes and Patient Experience Study', *Journal of Clinical Oncology*, vol. 31, no. 17, pp. 2136–2145.
- Smith S, Case L, Waterhouse K, Pettitt N, Beddard L, Oldham J, et al. 2012, *A blueprint of care for teenagers and young adults with cancer*, Teenage Cancer Trust, Manchester, UK.
- Steer B, Marx G, Singhal N, McJannett M, Goldstein D, Prowse R 2009, 'Cancer in older people: a tale of two disciplines', *Internal Medicine Journal*, vol. 39, pp. 771–775.
- Tan SY, Turner J, Kerin-Ayres K, Butler S, Deguchi C, Khatri S, et al. 2019, 'Health concerns of cancer survivors after primary anti-cancer treatment', *Support Cancer Care*, no. 10, pp. 3739–3747.
- Temel J, Greer J, Muzikansky A, Gallagher E, Admane S, Jackson V 2010, 'Early palliative care for patients with non-metastatic non-small cell lung cancer', *New England Journal of Medicine*, vol. 363, no. 8, pp. 733–742.
- Transplantation Society of Australia and New Zealand (TSANZ) 2024, 'Clinical guidelines for organ transplantation from deceased donors, Version 1.13', viewed 11 December 2024, <[https://tsanz.com.au/storage/Guidelines/TSANZ\\_Clinical\\_Guidelines\\_Version-113\\_250307.pdf](https://tsanz.com.au/storage/Guidelines/TSANZ_Clinical_Guidelines_Version-113_250307.pdf)>.
- Vardy JL, Raymond JC, Koczwara B, Lisy K, Cohn RJ, Joske D, et al. 2019, 'Clinical Oncology Society of Australia position statement on cancer survivorship care', *Australian Journal of General Practice*, vol. 48, no.12, pp. 833–836.
- Vijayvergia N, Denlinger CS 2015, 'Lifestyle factors in cancer survivorship: where we are and where we are headed', *Journal of Personalized Medicine*, vol. 5, no. 3, pp. 243–263.
- Vogel A, Bridgewater J, Edeline J, Kelley RK, Klumpen HJ, Malka D, et al. 2023, 'Biliary tract cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up'. *Annals of Oncology*, vol. 34, no. 2, pp. 127–140.
- Vogel A, Ducreux M, ESMO Guidelines Committee 2025, 'ESMO Clinical Practice Guideline interim update on the management of biliary tract cancer', *ESMO Open*, vol. 10, no. 1, p. 104003.
- Weber SM, Ribero D, O'Reilly EM, Kokudo N, Miyazaki M, Pawlik TM 2015, 'Intrahepatic Cholangiocarcinoma: expert consensus statement', *American International Hepato-Pancreato-Biliary Association*, no. 17, pp. 669–680.
- Wildiers H, Heeren P, Puts M, Topinkova E, Maryska LG, Janssen-Heijnen MLG, et al. 2014, 'International Society of Geriatric Oncology Consensus on geriatric assessment in older patients with cancer', *Journal of Clinical Oncology*, vol. 32, no. 24, pp. 2595–2603.
- World Health Organization (WHO) 2018, 'Malnutrition', WHO, Geneva, viewed 18 July 2019, <<https://www.who.int/news-room/fact-sheets/detail/malnutrition>>.
- Zimmermann C, Swami N, Krzyzanowska M, Hannon B, Leighl N, Oza A, et al. 2014, 'Early palliative care for patients with advanced cancer: a cluster-randomised controlled trial', *Lancet*, vol. 383, no. 9930, pp. 1721–1730.



