

# Understanding Pancreatic Cancer

*A guide for people with cancer, their families and friends*



For information & support, call **13 11 20**

## Understanding Pancreatic Cancer

A guide for people with cancer, their families and friends

First published May 1995. This edition March 2026.

© Cancer Council Australia 2026. ISBN 978 1 923073 46 3

*Understanding Pancreatic Cancer* is reviewed approximately every 3 years.

Check the publication date above to ensure this copy is up to date.

Editor: Nicole MacKee. Designer: Eleonora Pelosi. Printer: IVE Group.

### Acknowledgements

This edition has been developed by Cancer Council NSW on behalf of all other state and territory Cancer Councils as part of a National Cancer Information Subcommittee initiative. All updated content has been clinically reviewed by: Prof Lorraine Chantrill, Honorary Clinical Professor, University of Wollongong, and Head of Department, Medical Oncology, Illawarra Shoalhaven Local Health District, NSW. This edition is based on the previous edition, which was reviewed by the following panel: Prof Lorraine Chantrill (see above); Karen Baker, Consumer; Michelle Denham, 13 11 20 Consultant, Cancer Council WA; Prof Anthony J Gill, Surgical Pathologist, Royal North Shore Hospital and The University of Sydney, NSW; A/Prof Koroush Haghighi, Liver, Pancreas and Upper Gastrointestinal Surgeon, Prince of Wales and St Vincent's Hospitals, NSW; Dr Meredith Johnston, Radiation Oncologist, Liverpool and Campbelltown Hospitals, NSW; Dr Brett Knowles, Hepato-Pancreato-Biliary and General Surgeon, Royal Melbourne Hospital, Peter MacCallum Cancer Centre, and St Vincent's Hospital, VIC; Rachael Mackie, Upper GI – Clinical Nurse Consultant, Peter MacCallum Cancer Centre, VIC; Prof Jennifer Phillip, Chair of Palliative Care, University of Melbourne, and Palliative Medicine Physician, St Vincent's Hospital, Peter MacCallum Cancer Centre and Royal Melbourne Hospital, VIC; Lucy Pollerd, Social Worker, Peter MacCallum Cancer Centre, VIC; Rose Rocca, Senior Clinical Dietitian – Upper GI, Peter MacCallum Cancer Centre, VIC; Stefanie Simnadis, Clinical Dietitian, St John of God Subiaco Hospital, WA. We would like to thank all the health professionals, consumers and editorial teams who have worked on current and previous editions of this title.

### Note to reader

Always consult your doctor about matters that affect your health. This booklet is intended as a general introduction to the topic and should not be seen as a substitute for medical, legal or financial advice. You should obtain independent advice relevant to your specific situation from appropriate professionals, and you may wish to discuss issues raised in this booklet with them. All care is taken to ensure that the information in this booklet is accurate at the time of publication. Please note that information on cancer, including the diagnosis, treatment and prevention of cancer, is constantly being updated and revised by medical professionals and the research community. Cancer Council Australia and its members exclude all liability for any injury, loss or damage incurred by use of or reliance on the information provided in this booklet.

### Cancer Council

Cancer Council is Australia's peak non-government cancer control organisation. Through the 8 state and territory Cancer Councils, we provide a broad range of programs and services to help improve the quality of life of people living with cancer, their families and friends. Cancer Councils also invest heavily in research and prevention. To make a donation and help us beat cancer, visit [cancer.org.au](http://cancer.org.au) or call your local Cancer Council.



*Cancer Council acknowledges Traditional Custodians of Country throughout Australia and recognises the continuing connection to lands, waters and communities. We pay our respects to Aboriginal and Torres Strait Islander cultures and to Elders past and present.*



**Cancer Council Australia** Level 2, 320 Pitt Street, Sydney NSW 2000 ABN 91 130 793 725  
**Telephone** 02 8256 4100 **Email** [info@cancer.org.au](mailto:info@cancer.org.au) **Website** [cancer.org.au](http://cancer.org.au)

# About this booklet

This booklet has been prepared to help you understand more about pancreatic cancer.

Many people feel shocked and upset when told they have pancreatic cancer. We hope this booklet will help you, your family and friends understand how pancreatic cancer is diagnosed and treated. We also include information about support services.

We cannot give advice about the best treatment for you. You need to discuss this with your doctors. However, this information may answer some of your questions and help you think about what to ask your treatment team (see page 66 for a question checklist).

This booklet does not need to be read from cover to cover – just read the parts that are useful to you. Some medical terms that may be unfamiliar are explained in the glossary (see page 67). You may also like to pass this booklet to your family and friends for their information.

**How this booklet was developed** – This information was developed with help from a range of health professionals and people affected by pancreatic cancer. It is based on clinical practice guidelines for pancreatic cancer.<sup>1-2</sup>



If you or your family have any questions or concerns, call **Cancer Council 13 11 20**. We can send you more information and connect you with support services in your area. You can also visit your local Cancer Council website (see back cover).



# Contents

<b>What is cancer?</b>	<b>4</b>
<b>The pancreas</b>	<b>6</b>
<b>Key questions</b>	<b>8</b>
What is pancreatic cancer?	8
What are the main types?	8
What are the symptoms?	9
What are the risk factors?	10
Does pancreatic cancer run in families?	11
How common is pancreatic cancer?	11
What can I expect after diagnosis?	12
Where should I have treatment?	13
Which health professionals will I see?	14
<b>Diagnosis</b>	<b>16</b>
Blood tests	16
Imaging scans	17
Tissue sampling	20
Staging pancreatic cancer	23
Prognosis	25
<b>Making treatment decisions</b>	<b>27</b>
<b>Treatment to remove the cancer</b>	<b>29</b>
Treatments before or after surgery	29
Surgery to remove the cancer	30
What to expect after surgery	34
What if the cancer returns?	35

## Treatment to manage cancer and symptoms **38**

Surgery to relieve symptoms	39
Chemotherapy	41
Radiation therapy	43
How palliative care can help	46

## Managing your diet and nutrition **49**

Coping with the changes	49
Seeing a dietitian	50
Enzyme replacement therapy	51
Nausea and vomiting	54
Diarrhoea	56
Diabetes	57

## Looking after yourself **58**

## Living with pancreatic cancer **60**

When the cancer is advanced	61
-----------------------------	----

## Seeking support **62**

Support from Cancer Council	63
Useful websites	64

## Caring for someone with cancer **65**

## Question checklist **66**

## Glossary **67**

## How you can help **72**

## Key to icons

Icons are used throughout this booklet to indicate:



More information



Alert



Personal story



Tips

**Is this Cancer Council booklet helpful?**  
Please follow this QR code for a quick 3-minute survey, or call 13 11 20 to provide your feedback.



SCAN ME

# What is cancer?


Cancer is a disease of the cells. Cells are the body's basic building blocks - they make up tissues and organs. The body constantly makes new cells to help us grow, replace worn-out tissue and heal injuries.

Normally, cells multiply and die in an orderly way, so that each new cell replaces one lost. Sometimes, however, cells become abnormal and keep growing. These abnormal cells may turn into cancer.

In solid cancers, such as pancreatic cancer, the abnormal cells form a mass or lump called a tumour. In some cancers, such as leukaemia, the abnormal cells build up in the blood.

## How cancer starts

---





Not all tumours are cancer. Benign tumours tend to grow slowly and usually don't move into other parts of the body or turn into cancer. Cancerous tumours, also known as malignant tumours, have the potential to spread. They may invade nearby tissue, destroying normal cells. The cancer cells can break away and travel through the bloodstream or lymph vessels to other parts of the body.

The cancer that first develops in a tissue or organ is called the primary cancer. It is considered localised cancer if it has not spread to other parts of the body. If the primary cancer cells grow and form another tumour at a new site, it is called a secondary cancer or metastasis. A metastasis keeps the name of the original cancer. For example, pancreatic cancer that has spread to the liver is called metastatic pancreatic cancer, even though the main symptoms may be coming from the liver.

## How cancer spreads

### .....▶ Malignant cancer



# The pancreas

The pancreas is a long, flat gland that lies between your stomach and spine. It is about 13–15 cm long and is divided into 3 main parts:

- the large, rounded end, called the head of the pancreas
- the middle part, known as the body
- the narrow end, called the tail, that sits up next to your spleen.

A tube called the pancreatic duct connects the pancreas to the first part of the small bowel (duodenum). Another tube, called the common bile duct, runs through the head and joins with the pancreatic duct to connect the liver and gall bladder to the duodenum.

## What the pancreas does


The pancreas has 2 main jobs. It makes digestive juices (known as its exocrine function) and hormones (its endocrine function).

**Exocrine function** – The pancreas is part of the digestive system, which helps the body digest food and turn it into energy. Exocrine cells make pancreatic enzymes, which are digestive juices. The pancreatic duct carries these juices from the pancreas into the duodenum, where they help to break down food. Most of the pancreas is made up of exocrine tissue.

**Endocrine function** – The pancreas is also part of the endocrine system. This is a group of glands that makes the body's hormones. Endocrine cells in the pancreas make hormones that control blood sugar levels, the amount of acid produced by the stomach, and how quickly food is absorbed. For example, the hormone insulin decreases the level of sugar in the blood, while the hormone glucagon increases it.

# The pancreas in the body

---



# Key questions

## Q: What is pancreatic cancer?

**A:** Pancreatic cancer is cancer that starts in any part of the pancreas (see diagram previous page). About 3 in 4 pancreatic cancers are found in the head of the pancreas.

Pancreatic cancer can spread to nearby lymph nodes and to the lining of the abdomen (peritoneum). Cancer cells may also travel through the bloodstream to other parts of the body, such as the liver.

## Q: What are the main types?

**A:** There are 2 main groups of pancreatic cancer:

**Exocrine tumours** – These make up more than 90% of pancreatic cancers. The most common type is called adenocarcinoma, and it starts in the exocrine cells lining the pancreatic duct. Less common types include adenosquamous carcinoma, acinar cell carcinoma, squamous cell carcinoma and undifferentiated carcinoma.

**Neuroendocrine tumours (NETs)** – About 9% of cancers in the pancreas are pancreatic NETs. These start in the endocrine cells (see page 6).

This booklet is about exocrine tumours of the pancreas, particularly pancreatic adenocarcinomas. For more information about how pancreatic NETs are diagnosed and treated, see our *Understanding Neuroendocrine Tumours* fact sheet.

## Q: What are the symptoms?

**A:** Early-stage pancreatic cancer rarely causes obvious symptoms. Symptoms may not appear until the cancer is large enough to affect nearby organs or has spread.

The first symptom of pancreatic cancer is often jaundice. Signs of jaundice may include yellowish skin and eyes, dark urine, pale bowel motions and itchy skin. Jaundice is caused by the build-up of bilirubin, a dark yellow-brown substance found in bile. Bilirubin can build up if pancreatic cancer blocks the common bile duct.

Other common symptoms of pancreatic cancer include:

- appetite loss
- nausea (with or without vomiting)
- unexplained weight loss
- pain in the upper abdomen, side or back, which may cause you to wake up at night
- changed bowel motions – including diarrhoea, severe constipation, or pale, oily, foul-smelling stools (poo) that are difficult to flush away in the toilet
- bloating and passing wind and burping more than usual
- newly diagnosed type 2 diabetes
- fatigue (feeling very tired).

These symptoms can also occur in many other conditions and do not necessarily mean that you have cancer. If you are worried or have ongoing symptoms, speak with your general practitioner (GP).



For an overview of what to expect during all stages of your cancer care, visit [cancer.org.au/cancercareguides/pancreatic-cancer](https://cancer.org.au/cancercareguides/pancreatic-cancer). This is a short guide to what is recommended, from diagnosis to treatment and beyond.

*“I went to the doctor because I was itchy and had constant diarrhoea. My GP initially thought it was gallstones and sent me for routine tests. After the CT scan I went into hospital for a laparoscopy and then had a biopsy, which confirmed I had cancer.”* JAN

## Q: What are the risk factors?

**A:** The causes of pancreatic cancer are not known, but research has shown that people with certain risk factors are more likely to develop pancreatic cancer. Factors that are known to increase the risk of getting pancreatic cancer include:

- smoking tobacco (people who smoke are about twice as likely to develop pancreatic cancer as people who don't smoke)
- obesity
- ageing
- long-term diabetes (but diabetes can also be caused by the pancreatic cancer, see page 57)
- long-term pancreatitis (inflammation of the pancreas)
- certain types of cysts in the pancreatic duct known as intraductal papillary mucinous neoplasms (IPMNs) – these should be assessed by an appropriate specialist
- stomach infections caused by the *Helicobacter pylori* bacteria (which can also cause stomach ulcers)
- family history and inherited conditions (see opposite page)
- heavy exposure to certain chemicals used in some workplaces.

Having risk factors does not mean you will definitely get cancer, but talk to your doctor if you are concerned. Some people with pancreatic cancer have no known risk factors.

## **Q: Does pancreatic cancer run in families?**

**A:** Most people diagnosed with pancreatic cancer do not have a family history of the disease. Only about 5–10% of people who develop pancreatic cancer have inherited a faulty gene that increases the risk of developing pancreatic cancer.

You may have inherited a faulty gene linked to pancreatic cancer if:

- 2 or more of your close family members (such as a parent or sibling) have had pancreatic cancer
- there is a family history of a genetic condition, such as Peutz-Jeghers syndrome, the familial breast cancer genes (BRCA1 and BRCA2), familial atypical multiple mole melanoma (FAMMM) syndrome, Lynch syndrome and hereditary pancreatitis.

Genetic testing aims to find inherited faulty genes that may increase a person's risk of developing some cancers. People with a strong family history of cancer can go to a family cancer clinic for genetic counselling and tests.

For more information about family history and pancreatic cancer, talk to your doctor or local family cancer clinic.

## **Q: How common is pancreatic cancer?**

**A:** About 4600 people are diagnosed with pancreatic cancer in Australia each year. More than 80% of those diagnosed are over the age of 60.

Pancreatic cancer is about the 8th most common cancer diagnosed in Australia and it affects men and women at about the same rate.<sup>3</sup>

## **Q: What can I expect after diagnosis?**

**A:** It's common to have many questions and concerns about what a diagnosis of pancreatic cancer will mean for you. This booklet has chapters that explain the process of diagnosis, treatment options for removing the cancer and managing symptoms, and coping with physical changes that affect what you can eat.

### **Diagnosis stage (pages 16–26)**

You will have various tests to confirm that you have pancreatic cancer, and to work out which type and how far it has progressed. The results will help guide decisions about treatment.

### **Treatment to remove the cancer (pages 29–37)**

About 20% of people with pancreatic cancer can have surgery to remove the cancer. They may also be offered other treatments before or after the surgery.

### **Treatment to manage cancer and symptoms (pages 38–48)**

Most people with pancreatic cancer are diagnosed at a stage when the cancer cannot be removed. Treatment will aim to control the cancer, manage symptoms and improve quality of life.

### **Managing your diet and nutrition (pages 49–57)**

People with pancreatic cancer often need to adapt to changes in how their body processes food, either because of the cancer itself or because of the treatment.

If you are feeling overwhelmed or would just like to talk through your concerns, you can call Cancer Council 13 11 20 to speak to one of our experienced health professionals. You can also listen to our podcast *The Thing About Cancer* at [cancercouncil.com.au/podcasts](http://cancercouncil.com.au/podcasts).

## Q: Where should I have treatment?

**A:** Treatment for pancreatic cancer is highly specialised. This is particularly the case for the surgery for early pancreatic cancer, such as the Whipple procedure (see pages 32–33).

There is strong evidence that outcomes are better when people have their treatment in a specialist centre that sees a lot of people with pancreatic cancer. These high-volume centres have multidisciplinary teams of health professionals experienced in treating pancreatic cancer (see the next 2 pages).

Specialist centres offer access to a wide range of treatment options – including treatments available only through clinical trials (see page 28) – but you may need to travel away from home for treatment.

Sometimes the multidisciplinary team from a specialist centre will be able to advise your local specialist. You may find that you can visit the specialist centre to confirm the diagnosis and work out a treatment plan to have much of your treatment closer to your home.

To find a treatment centre that specialises in pancreatic cancer, talk to your GP.



If you live in a rural or regional area and have to travel a long way for appointments or treatment, you may be able to get financial assistance towards the cost of accommodation or travel. To check whether you are eligible or to apply for this assistance, speak to your GP or the hospital social worker, or call Cancer Council 13 11 20.

## Q: Which health professionals will I see?

**A:** Your GP will arrange the first tests to assess your symptoms. If these tests do not rule out cancer, you will usually be referred to a specialist, such as a gastroenterologist or surgeon. The specialist will arrange further tests. If pancreatic cancer is diagnosed,

### Health professionals you may see

<b>pancreatic or HPB (hepato-pancreato-biliary) surgeon</b>	operates on the liver, bile ducts, pancreas and surrounding organs
<b>gastroenterologist</b>	diagnoses and treats disorders of the digestive system; may diagnose pancreatic cancer, perform endoscopy, and insert stents to clear blocked bile ducts
<b>medical oncologist</b>	treats cancer with drug therapies such as chemotherapy, immunotherapy and targeted therapy (systemic treatment)
<b>radiation oncologist</b>	treats cancer by prescribing and overseeing a course of radiation therapy
<b>endocrinologist</b>	diagnoses, treats and manages hormonal disorders, including diabetes
<b>radiologist</b>	analyses x-rays and scans; an interventional radiologist may also perform a biopsy under ultrasound or CT, and deliver some treatments
<b>cancer care coordinator</b>	coordinates your care, liaises with other members of the MDT and supports you and your family throughout treatment; care may also be coordinated by a clinical nurse consultant (CNC) or clinical nurse specialist (CNS)

the specialist will consider treatment options. Often these will be discussed with other health professionals at what is known as a multidisciplinary team (MDT) meeting. During and after treatment, you will see a range of health professionals who specialise in different aspects of your care.

<b>nurse</b>	administers drugs and provides care, information and support throughout treatment
<b>palliative care team</b>	a team of specialist doctors, nurses and allied health workers who work closely with the GP and cancer team to help control symptoms and maintain quality of life
<b>dietitian</b>	helps with nutrition concerns and recommends changes to diet during treatment and recovery
<b>social worker</b>	links you to support services and helps you with emotional, practical and financial issues
<b>psychologist, counsellor</b>	help you manage your emotional response to diagnosis and treatment
<b>physiotherapist, exercise physiologist</b>	help restore movement and mobility, and improve fitness and wellbeing
<b>occupational therapist</b>	assists in adapting your living and working environment to help you resume usual activities after treatment

# Diagnosis

If your doctor thinks you may have pancreatic cancer, you will need some tests to confirm the diagnosis. These may include blood tests, CT, MRI and other imaging scans, endoscopic tests and tissue sampling (called a biopsy). You may also have tests to check for gene changes in the cancer. The tests you have will depend on the symptoms, type and stage of the cancer. You will not have all the tests described in this chapter.

## Blood tests

You are likely to have blood tests to check your general health and see how well your liver and kidneys are working. Some blood tests look for proteins produced by cancer cells. These proteins are known as tumour markers.

## Tumour markers

Many people with pancreatic cancer have higher levels of particular tumour markers in their bloodstream. The markers that doctors will look for are CA 19-9 (carbohydrate antigen) and CEA (carcinoembryonic antigen). On their own, however, these tumour markers can't be used to diagnose pancreatic cancer. This is because some people with pancreatic cancer have normal levels of these markers, and other conditions can also raise the levels of these markers in the bloodstream.

It is normal for the levels of these tumour markers to go up and down a little. Your doctor will look for sharp increases and overall patterns. Raised levels can tell your doctor more about the cancer and, after diagnosis, can also show how well the treatment is working.

## Imaging scans

Imaging scans are tests that create pictures of the inside of the body. Different scans can provide different details about the cancer.

You will usually have at least one of the following scans during diagnosis and treatment.

### CT scan

Most people suspected of having pancreatic cancer will have a CT (computerised tomography) scan. This scan uses x-ray beams to take many pictures of the inside of your body and then compiles them into one detailed, cross-sectional picture.

A CT scan is usually done at a hospital or a radiology clinic. Before the scan, a liquid dye (called contrast) will be injected into a vein to help make the pictures clearer. The dye travels through your bloodstream to the pancreas and nearby organs and helps show up any abnormal areas. This may cause you to feel hot all over, may give you a strange taste in your mouth and you could feel as if you need to pass urine (pee). These reactions are temporary and usually go away in a few minutes, but tell your treatment team if you feel unwell.

The CT scanner is large and round like a doughnut. You will need to lie still on an examination table while the scanner moves around you. The scan itself is painless and takes only a few minutes, but getting it set up can take 10–30 minutes.



Before having scans, tell the doctor if you have any allergies or have had a reaction to dyes during previous scans. You should also let them know if you have diabetes or kidney disease, or are pregnant or breastfeeding.

## Endoscopic tests

Endoscopic tests can show blockages or inflammation in the common bile duct, stomach and duodenum. For these tests, you will have an endoscopy, a procedure that is usually done as day surgery by a specialist doctor called a gastroenterologist. The doctor passes a long, flexible tube with a light and small camera on the end (endoscope) down your throat into your digestive tract. There are 2 main types of endoscopic tests:

**Endoscopic ultrasound (EUS)** – An EUS uses an endoscope with an ultrasound probe (transducer) attached. The endoscope is passed through your mouth into the small bowel. The transducer makes soundwaves that create detailed pictures of the pancreas and ducts. This helps to locate small tumours and shows if the cancer has spread into nearby tissue. A biopsy of the cancer can be taken at the same time (see page 20).

**Endoscopic retrograde cholangiopancreatography (ERCP)** – An ERCP is used to take an x-ray of the common bile duct and pancreatic duct. The doctor uses the endoscope to guide a tube into the bile duct and insert a small amount of dye. The x-ray images show blockages or narrowing that might be caused by cancer. ERCP may also be used to put a thin plastic or metal tube (stent) into the bile duct to keep it open (see page 40).

During an endoscopic procedure, the doctor can also take a tissue or fluid sample (biopsy) to help with the diagnosis (see pages 20–21).

You will be asked not to eat or drink (fast) for 6 hours before an endoscopic test. The doctor will give you medicine to help you relax and feel as comfortable as possible. Because of this medicine, you shouldn't drive or operate machinery until the next day.

Having an endoscopic test has some risks, including infection, bleeding and inflammation of the pancreas (pancreatitis). These complications are not common. Your doctor will explain the risks before asking you to agree (consent) to the procedure.

## **MRI and MRCP scans**

In some cases, you may also have another type of scan such as an MRI or MRCP scan. An MRI (magnetic resonance imaging) scan uses a powerful magnet and radio waves to create detailed cross-sectional pictures of the pancreas and nearby organs. An MRCP (magnetic resonance cholangiopancreatography) scan is a different type of MRI scan that produces more detailed images and can be used to check the common bile duct for a blockage (obstruction).

An MRI or MRCP takes about an hour and you will be able to go home when it is over. Before the scan, you may be asked not to eat or drink (fast) for a few hours. You may also be given an injection of dye (contrast) to highlight the organs in your body.

During the scan, you will lie on a treatment table that slides into a large metal tube that is open at both ends. The noisy, narrow machine makes some people feel anxious or claustrophobic. If you think you might be distressed, mention this beforehand to your doctor or nurse. You may be given medicine to help you relax, and you will usually be offered headphones or earplugs. Also let the doctor or nurse know if you have a pacemaker or any other metallic object in your body, as this can interfere with the scan.

MRIs for pancreatic cancer are not always covered by Medicare. If this test is recommended, check with your treatment team what you may have to pay.

## PET–CT scan

Doctors sometimes use a PET (positron emission tomography) scan combined with a CT scan to help work out if the pancreatic cancer has spread or how it is responding to treatment.

It may take several hours to prepare for and complete a PET–CT scan. Before the scan you will be injected with a small amount of radioactive material, usually a glucose solution called fluorodeoxyglucose (FDG). Some cancer cells will show up brighter on the scan because they take up more of this solution than normal cells do.

PET–CT scans are specialised tests. They are not available in every hospital and may not be covered by Medicare, so talk to your medical team for more information.

## Tissue sampling

If imaging scans show there is a tumour in the pancreas, your doctor may remove a sample of cells or tissue from the tumour (biopsy).

This is the main way to confirm if the tumour is cancer and to work out exactly what type of cancer it is. A specialist doctor called a pathologist will examine the sample under a microscope to check for signs of cancer. Sometimes, the results are not clear and a second biopsy is needed.

A biopsy can be taken in different ways, including:

**With a needle** – A sample of cells may be collected with a fine needle (fine needle biopsy), or a tissue sample may be collected with a larger needle (core biopsy). A fine needle or core biopsy can be done during an endoscopic procedure (see pages 18–19). Another method is to insert

the needle through the skin of the abdomen, using an ultrasound or CT scan for guidance. You will be awake during the procedure, but you will be given a local anaesthetic so you do not feel any pain.

**During keyhole surgery** – Also called a laparoscopy or minimally invasive surgery, keyhole surgery is sometimes used to look inside the abdomen to see if the cancer has spread to other parts of the body. It can also be done to take tissue samples before any further surgery.

Keyhole surgery is done under general anaesthetic, so you will be asked not to eat or drink (fast) for 6 hours beforehand. If you take blood-thinning medicines or have diabetes, let your doctor or nurse know before the surgery as they may need to adjust your medicines in the days leading up to the procedure.

The procedure uses an instrument called a laparoscope (a long tube with a light and camera on the end). The camera projects images onto a screen so the doctor can see inside your body. The doctor will guide the laparoscope through a small cut near your bellybutton. The doctor can then insert other instruments through other small cuts to take the biopsy.

You will have stitches where the cuts were made. You may feel sore while you heal, so you will be given pain medicine during and after the operation, and to take at home. There is a small risk of infection or damage to an organ with a laparoscopy. Your doctor will explain the risks before asking you to agree to the procedure.

**During surgery to remove the tumour** – If you are having a larger operation to remove the tumour (see pages 30–33), your surgeon may take the tissue sample at that time.

## Genomic and genetic testing

---

Each cell in the human body has about 20,000 genes, which tell the cell what to do and when to grow and divide. Cancer starts because of changes to the genes (known as mutations).

Some people are born with a gene change that increases their risk of cancer (an inherited faulty gene), but most gene changes that cause cancer build up during a person's lifetime (acquired gene changes).

In some circumstances, your doctors may recommend extra tests to look for acquired gene changes (genomic or molecular tests) or inherited gene changes (genetic tests).

### Genomic testing

If you have pancreatic cancer, you may be offered extra tests on the biopsy sample known as genomic or molecular testing. This looks for gene changes and other features in the cancer cells that may help your doctors decide which treatments to recommend.

At the time of publication (March 2026), Medicare does not cover genomic testing for pancreatic cancer. These tests can be

expensive, so check what costs are involved and how helpful it would be. If you are having genomic or molecular testing as part of a research project, the costs may be covered.

### Genetic testing

Your doctor may suspect you have developed pancreatic cancer because you have inherited a faulty gene (e.g. because other members of your family have also had pancreatic cancer). In this case, they may refer you to a family cancer clinic for genetic counselling and extra tests.

These tests are known as genetic or germline tests. The results may help your doctor work out what treatment to recommend and can also provide important information for your blood relatives.

Genetic counselling can help you understand what tests are available to you and what the results mean for you and your family.

Medicare may cover the costs of genetic tests or you may need to pay for them – check this with your treatment team.

## Staging pancreatic cancer

The test results will show what type of pancreatic cancer it is, where in the pancreas it is, and whether it has spread. This is called staging, and it helps your doctors work out the best treatment options for your situation.

Pancreatic cancer is commonly staged using the TNM system, with each letter given a number that shows how advanced the cancer is:

- T stands for Tumour and refers to the size of the tumour or how close it is to major blood vessels. It will be given a score of T0–4. The higher the number, the more advanced the cancer is.
- N stands for Nodes and refers to whether the cancer has spread to lymph nodes. N0 means the cancer has not spread to nearby lymph nodes; N1 means the cancer has spread to 1–3 nodes, N2 means the cancer has spread to 4 or more lymph nodes.
- M stands for Metastasis. M0 means the cancer has not spread to other parts of the body; M1 means it has.

The TNM scores are combined to work out the overall stage of the cancer, from stage 1 to stage 4 (see the table on the next page). If you need help to understand staging, ask someone in your treatment team to explain it in a way that makes sense to you. You can also call Cancer Council 13 11 20.

*“I asked the surgeon what caused it, and he said, ‘We don’t know, Phil.’ He got me to focus on what had to be done and to just get on with it. It’s easy to say now that you need to get on the front foot and work with your treatment team, but the diagnosis is a terrific blow.”* PHIL

## Stages of pancreatic cancer

<p><b>stage 1</b></p>	<p>Cancer is small and found only in the pancreas.</p>	<p>Stage 1–2 cancers are considered early pancreatic cancer. They may also be called resectable, which means surgery to remove the cancer (see pages 30–33) may be an option if you are well enough. About 1 in 5 pancreatic cancers are stage 1–2 when first diagnosed.</p>
<p><b>stage 2</b></p>	<p>Cancer is large but has not spread outside the pancreas; or it is small and has spread to a few nearby lymph nodes.</p>	
<p><b>stage 3</b></p>	<p>Cancer has grown into nearby major blood vessels or into a lot of nearby lymph nodes.</p>	<p>Some stage 3 cancers are borderline resectable cancers, which means surgery to remove the cancer may be an option if other treatment can shrink the cancer first (see page 29). Other stage 3 cancers are called locally advanced, which means that surgery cannot remove the cancer, but treatments can relieve symptoms (see pages 38–48). About 30% of pancreatic cancers are stage 3 when first diagnosed.</p>
<p><b>stage 4</b></p>	<p>The cancer has spread to more distant parts of the body, such as the liver, lungs or lining of the abdomen. There may or may not be cancer in the lymph nodes.</p>	<p>This is called metastatic cancer. Surgery cannot remove the cancer, but treatments can relieve symptoms. About half of all pancreatic cancers are stage 4 when first diagnosed.</p>

## Prognosis

Prognosis means the expected outcome of a disease. You may wish to discuss your prognosis with your doctor, but it is not possible for anyone to predict the exact course of the disease.

To work out your prognosis, your doctor will consider:

- test results
- the type, stage and location of the cancer
- how the cancer responds to initial treatment
- your medical history
- your age and general health.

As symptoms can be vague or go unnoticed, most pancreatic cancers are not found until they are advanced, which usually means treatment cannot remove all the cancer. If the cancer is diagnosed at an early stage and can be surgically removed, the prognosis may be better.

Discussing your prognosis and thinking about the future can be challenging and stressful. It is important to know that although the statistics for pancreatic cancer can be frightening, they are an average and may not apply to your situation. Talk to your doctor about how to interpret any statistics that you come across.

When pancreatic cancer is advanced, treatment will usually aim to control the cancer for as long as possible, relieve symptoms and improve quality of life. This is known as palliative treatment (see pages 38–48 for more information).

## Key points about diagnosing pancreatic cancer

---

### Main tests

- Blood tests can check your general health and look for tumour markers.
- CT scans help show where the cancer is in the pancreas, and if it has spread to other areas of the body.
- Endoscopic tests include EUS (endoscopic ultrasound) and ERCP (endoscopic retrograde cholangiopancreatography). A long, flexible tube with a light and tiny camera is passed into your digestive system. The camera shows the pancreas and ducts, and a tissue sample can be taken.
- A tissue sample may be removed during a scan or procedure. This is called a biopsy. The sample will be examined under a microscope to see whether cancer is present and, if so, what type it is.

---

### Other tests

- An MRI scan is sometimes used to help diagnose pancreatic cancer. Some people have a type of MRI scan called an MRCP that helps show blockages in the common bile duct.
- Some people have a PET-CT scan to see if pancreatic cancer has spread.
- You may be referred to a family cancer clinic for genetic tests if your doctor suspects that the cancer is related to an inherited faulty gene.

---

### Staging

- The stage shows how far the cancer has spread throughout the body. Most pancreatic cancers are not found until they are advanced.
  - Staging helps your treatment team decide on suitable treatment options.
-



# Making treatment decisions

Sometimes it is difficult to decide on the type of treatment to have. You may feel that everything is happening too fast, or you might be anxious to get started.

Check with your specialist how soon treatment should begin, as it may not affect the success of the treatment to wait a short time. Ask them to explain the options, and take what time you can before making a decision.

**Know your options** – Understanding the disease, the available treatments, possible side effects and any extra costs can help you weigh up the options and make a well-informed decision. Check if the specialist is part of a multidisciplinary team (see page 15) and if the treatment centre is the most appropriate one for you – you may be able to have treatment closer to home, or it might be worth travelling to a centre that specialises in a particular treatment.

**Record the details** – When your doctor first says you have cancer, you may not remember everything you are told. Taking notes can help. If you would like to record the discussion, ask your doctor first. It is a good idea to have a family member or friend go with you to appointments to join in the discussion, write notes or simply listen.

**Ask questions** – If you are confused or want to check anything, it is important to ask your specialist questions. Try to prepare a list before appointments (see page 66 for suggestions). If you have a lot of questions, you could talk to a cancer care coordinator or nurse.

**Consider a second opinion** – You may want to get a second opinion from another specialist to confirm or clarify your specialist's recommendations or reassure you that you have explored all of your options. Specialists are used to people doing this. Your GP or specialist can refer you to another specialist and send your initial results to that person. You can get a second opinion even if you have started treatment or still want to be treated by your first doctor. You might decide you would prefer to be treated by the second specialist.

**It's your decision** – Adults have the right to accept or refuse any treatment they are offered. For example, some people with advanced cancer choose treatment that has significant side effects even if it gives only a small benefit for a short period of time. Others decide to focus their treatment on quality of life. You may want to discuss your decision with the treatment team, GP, family and friends.

## Should I join a clinical trial?

---

Your doctor or nurse may suggest you take part in a clinical trial. Doctors run clinical trials to test new or modified treatments and ways of diagnosing disease to see if they are better than current methods. For example, if you join a randomised trial for a new treatment, you will be chosen at random to receive either the best existing treatment or the modified new treatment. Over the years, trials have improved treatments and

led to better outcomes for people diagnosed with cancer.

You may find it helpful to talk to your specialist, clinical trials nurse or GP, or to get a second opinion. If you decide to take part in a clinical trial, you can withdraw at any time. For more information, visit [australiancancertrials.gov.au](http://australiancancertrials.gov.au).

► See our *Understanding Clinical Trials and Research* booklet.

# Treatment to remove the cancer

This chapter gives an overview of treatments used for many stage 1-2 (early) pancreatic cancers and some stage 3 pancreatic cancers. You can read more about the stages of pancreatic cancer on pages 23-24.

The treatment options described in this chapter will be suitable for only about 1 in 5 people with pancreatic cancer, as most people are diagnosed at a later stage. Surgery to remove the cancer, in combination with chemotherapy and possibly radiation therapy, is generally the most effective treatment for early-stage pancreatic cancer. It is important that the surgery is done by a surgeon who is part of a multidisciplinary team in a specialist pancreatic cancer treatment centre (see page 15).

## Treatments before or after surgery

Your surgeon may recommend other treatments before surgery to shrink the tumour (called neoadjuvant therapies), or after surgery to destroy any remaining cancer cells (called adjuvant therapies). They may include:

- **chemotherapy** – the use of drugs to kill or slow the growth of cancer cells (see pages 41-42)
- **chemoradiation** – chemotherapy combined with radiation therapy (see pages 43-44).



Your health care team may talk to you about doing some exercises to help improve your fitness before surgery. This may be called prehabilitation. Ask your treatment team about prehabilitation and what exercises would be best for you.

## Surgery to remove the cancer

Surgery to remove the tumour (resection) is the most common treatment for people with early-stage cancer who are in good health. It may also be considered for some stage 3 cancers, usually with chemotherapy (and sometimes radiation therapy) to shrink the tumour first. These stage 3 cancers are known as borderline resectable cancers, which means that surgery alone may or may not be able to remove all of the tumour.

The aim of resection is to remove all the tumour from the pancreas, as well as a margin of healthy tissue. The type of surgery you have will depend on the size and location of the tumour, your general health and your preferences. Your surgeon will talk to you about the most appropriate surgery for you, as well as the risks and any possible complications. Types of surgery include:

**Whipple procedure** – This treats tumours in the head of the pancreas. Also known as pancreaticoduodenectomy, it is the most common surgery for pancreatic tumours. See pages 32–33 for more information.

**Distal pancreatectomy** – The surgeon removes only the tail of the pancreas, or the tail and a portion of the body of the pancreas. The spleen is usually removed as well. The spleen helps the body fight infections, so if it is removed you are at higher risk of some types of bacterial infection. Your doctor may recommend vaccinations before and after a distal pancreatectomy.

**Total pancreatectomy** – When cancer is large or there are many tumours, the entire pancreas and spleen may be removed, along with the gallbladder, common bile duct, part of the stomach and small bowel, and nearby lymph nodes.

## If the cancer has spread

During surgery to remove the cancer, the surgeon may find that the cancer has spread around one or more of the major blood vessels in the area or into the lining of the abdomen (peritoneum). This may occur even if you had several scans and tests beforehand.

If this happens, the surgeon will not be able to remove the cancer. However, they may be able to perform procedures (such as a bypass) to relieve some of the symptoms caused by the cancer. See pages 39–40.

## How the surgery is done

---

Surgery for pancreatic cancer is carried out in hospital under a general anaesthetic. It is difficult surgery and should only be performed by surgeons with a lot of experience in doing this operation. There are 3 main approaches:

**Open surgery** – This involves one larger cut in the abdomen so the surgeon can remove the cancer.

**Keyhole surgery** – This involves several small cuts in the abdomen (belly). It is sometimes called laparoscopic or minimally invasive surgery. The surgeon inserts a long, thin instrument with a light and camera (laparoscope) into one of the cuts and uses images from the camera for guidance.

The surgeon inserts tools into the other cuts to remove the cancer.

**Robotic-assisted surgery** – This is a type of keyhole surgery. The surgeon sits at a control panel to see a 3-dimensional image and moves robotic arms that hold the instruments.

Open surgery is usually the best approach for pancreatic cancer, but keyhole or robotic-assisted surgery may be an option in some circumstances. Talk to your surgeon about what options are available to you. Ask about the risks and benefits of each approach, and check if there are any extra costs.


► See our *Understanding Surgery* booklet for more details.

## Having a Whipple procedure

The Whipple procedure (pancreaticoduodenectomy) is a major, complex operation. It has to be done by a specialised pancreatic surgeon or a hepato-pancreato-biliary (HPB) surgeon.

The surgeon removes the part of the pancreas with the cancer (usually the head); the first part of the small bowel (duodenum); part of the stomach; the gallbladder; and part of the common bile duct.

### Before



A Whipple procedure is a long operation. It usually lasts 5–8 hours.




As your surgeon will explain, this surgery is complex and there is a chance of serious problems, such as major bleeding or leaking where the remaining organs have been reconnected.

Then the surgeon reconnects the remaining part of the pancreas, common bile duct and stomach (or duodenum) to different sections of the small bowel to keep the digestive tract working.

Once these organs are reconnected, food, pancreatic juices and bile can continue to flow into the small bowel for the next stage of digestion. Many people need to change their diet after a Whipple procedure.

### After



Most people stay in hospital for 1–2 weeks after surgery, and full recovery takes at least 8–12 weeks. Your team will encourage you to move around and start gentle exercise as soon as you are ready.



For tips on adjusting to issues with food after a Whipple procedure, see the *Managing your diet and nutrition* chapter (pages 49–57).

## What to expect after surgery

While you are recovering after surgery, your health care team will check your progress and help you with the following:

**Pain control** – You will have some pain and discomfort for several days after surgery. You will be given pain medicines to manage this. If you are in pain when you return home, talk to your doctors about a prescription for pain medicine.

**Surgical drain** – You may have a thin tube placed in the abdomen to drain fluid into a small bag or bottle. The fluid can then be checked for potential problems. The tube is usually removed after a few days but may be left in for longer. Surgical drains are never permanent.

**Drips and tubes** – While in hospital, you will have a drip to replace your body's fluids. At first, you may not be able to eat or drink (nil by mouth). You'll then be on a liquid diet before slowly returning to normal food. A temporary feeding tube may be put into the small bowel during the operation. This tube provides extra nutrition until you can eat and drink normally again. The hospital dietitian can help you manage changes to eating.

**Enzyme supplements** – Many people will need to take tablets known as pancreatic enzyme replacement therapy (PERT) after surgery (see page 51). These are taken with each meal to help digest fat and protein.

**Insulin therapy** – Because the pancreas produces insulin, people who have had all or some of their pancreas removed may develop diabetes after surgery and need regular insulin injections (up to 4 times per day). A specialist doctor called an endocrinologist will help you develop a plan for managing diabetes. See page 57 for tips.

**Moving around** – Your health care team will probably encourage you to walk the day after surgery. They will also provide advice about when you can get back to your usual activity levels.

**Length of hospital stay** – Most people go home within 2 weeks, but if there are problems, you may need to stay in hospital longer.

You may need rehabilitation, which is a program to help you recover and regain physical strength, and adapt to changes, after surgery. This may be as an inpatient in a rehabilitation centre or through a home-based rehabilitation program.

## **What if the cancer returns?**

If the surgery successfully removes all of the cancer, you will have regular appointments to monitor your health, manage any long-term side effects and check that the cancer hasn't come back or spread.

Check-ups will become less frequent if you have no further problems. Between appointments, let your doctor know immediately of any symptoms or health problems.

Unfortunately, pancreatic cancer is difficult to treat and it may come back after treatment. This is known as a recurrence. Most of the time, surgery is not an option if you have a recurrence. Your doctors may recommend other types of treatment with the aim of reducing symptoms and improving quality of life.

The next chapter describes some of these treatments. You may also be able to get new treatments by joining a clinical trial (see page 28).



## Karen's story

*I had lost a lot of weight and started to pass dark urine, but I didn't think a lot of it until later when I went to the doctor and mentioned back pain. I was very lucky as she felt my stomach area and sent me for a CT scan, and they found a 5 cm tumour.*

*Pancreatic is not as common as other cancers and rarely mentioned in the media. But there have been improvements in surgery and it is treatable if caught in the early stages.*

*After my diagnosis, I had the Whipple procedure followed by chemotherapy and radiation.*

*It took about a year to really get over it. I still have trouble with digestion and have to take Creon tablets when eating.*

*After each of my radiation treatments, I would become very emotional. When I first started the radiation therapy, I burst into tears; it was such an imposing machine and I found the whole process overwhelming and a bit scary.*

*Before the pancreatic cancer diagnosis, I was a busy teacher and my children were all still at home. I now have the time to walk regularly and cancer was a great motivator for doing strength exercises. I've taken up my art again and attend an art group, which I find very therapeutic.*

*Because of the nature of my cancer, I would rate the fear of recurrence as 10 out of 10. I've already been told by the surgeon that I will have a shortened life span.*

*When you have a cancer diagnosis, you don't stop being a wife, a mother and friend, and some people don't stop working. There's still stuff in your life to get on with, and I find talking with a counsellor has helped me with the things that I can't talk to anyone else about.*

***“I've taken up my art again and attend an art group, which I find very therapeutic.”***

## Key points about treatment to remove the cancer

---

### The main treatment

- The main treatment for stage 1–2 pancreatic cancers and some stage 3 pancreatic cancers is surgery to remove the tumour (resection).
  - This is an option for about 1 in 5 people with pancreatic cancer.
- 

### Types of surgery

- A Whipple procedure treats tumours in the head of the pancreas. It removes the gallbladder and parts of the pancreas, small bowel (duodenum), common bile duct and stomach. Full recovery can take at least 8–12 weeks.
  - A distal pancreatectomy is sometimes used to treat pancreatic cancer found in the tail or body of the pancreas. It removes part of the pancreas and often the spleen.
  - In a total pancreatectomy, the entire pancreas and spleen are removed, along with the gallbladder, common bile duct, part of the stomach and small bowel, and nearby lymph nodes.
- 

### Other treatment options

- You may have chemotherapy, alone or in combination with radiation therapy, before or after surgery.
  - After part or all of the pancreas is removed, you may need to take pancreatic enzymes to help digest fat and protein, and have insulin injections if you develop diabetes.
  - Whatever treatment is needed, your health care team will be working to ensure your symptoms are well managed.
-



# Treatment to manage cancer and symptoms

Pancreatic cancer usually has no symptoms in its early stages, so many people are diagnosed when the cancer is advanced. If the cancer involves nearby organs or blood vessels (locally advanced – some stage 3 cancers) or has spread to other parts of the body (metastasised – all stage 4 cancers), surgery to remove the cancer may not be possible. Instead, treatments will focus on shrinking or slowing the growth of the cancer and relieving symptoms without trying to cure the disease. This is called palliative treatment.

Some people think that palliative treatment is only offered at the end of life. However, it can help at any stage of pancreatic cancer. It does not mean giving up hope – rather, it is about managing symptoms and living as fully and comfortably as possible. Palliative treatment is one aspect of palliative care (see pages 46–47). Many studies have shown that when palliative care is delivered early it can improve people’s quality of life.

Palliative treatments may include surgery, chemotherapy and radiation therapy, either on their own or in combination. These treatments may be used to help manage the cancer and relieve some common symptoms of advanced pancreatic cancer, such as:

- jaundice caused by narrowing of the common bile duct
- ongoing vomiting and weight loss caused by a blockage in the stomach or small bowel
- pain in the abdomen (belly) and middle back.

Many people with advanced pancreatic cancer have digestive problems. For example, a blockage in the pancreatic duct can stop the flow of the

digestive enzymes required to break down food. This can be treated with pancreatic enzyme supplements. See pages 49–57 for ways to change your diet to help manage some common problems.

## Surgery to relieve symptoms

If the tumour is pressing on the common bile duct, it can cause a blockage and prevent bile from passing into the bowel. Bile builds up in the blood, causing symptoms of jaundice, such as yellowing of the skin and whites of the eyes; itchy skin; reduced appetite, poor digestion and weight loss; dark urine; and pale stools. If cancer blocks the duodenum (first part of the small bowel), food cannot pass into the bowel and builds up in your stomach, causing nausea and vomiting.


Blockages of the common bile duct or duodenum are known as obstructions. Surgical options for managing these may include:

- **stenting** – inserting a small tube into the bile duct or duodenum (this is the most common method, see next page)
- **bypass surgery** – connecting the small bowel to the bile duct or gallbladder to redirect the bile around the blockage, and connecting a part of the bowel to the stomach to bypass the duodenum so the stomach can empty properly
- **gastroenterostomy** – connecting the stomach to the jejunum (middle section of the small bowel)
- **venting gastrostomy** – connecting the stomach to an opening in the abdomen so waste can be collected in a small bag outside the body.

Sometimes the surgeon may have planned to remove a pancreatic tumour but discovers during the surgery that the cancer has spread. If the tumour cannot be removed, the surgeon may perform one of the operations listed above to relieve symptoms.

## Inserting a stent

---



If the cancer cannot be removed and is pressing on the common bile duct or duodenum, you may need a stent. A stent is a small tube made of either plastic or metal. It holds the bile duct or duodenum open, letting the bile or food flow into the bowel again.

A bile duct stent is also known as a biliary stent. It is usually inserted using an endoscope passed through the mouth, stomach and duodenum until it reaches the bile duct. You may have this procedure as an outpatient or stay in hospital for 1–2 days.

Sometimes the stent needs to be inserted directly through the skin and liver into the bile duct. This is called a PTC (percutaneous transhepatic cholangiogram).

A duodenum stent is also known as a duodenal stent. It is usually inserted via the mouth using an endoscope.

Symptoms caused by the blockage usually go away over 2–3 weeks. Your appetite is likely to improve and you may gain some weight.

## Chemotherapy

Chemotherapy uses drugs to kill or slow the growth of cancer cells. It is used for pancreatic cancer at various stages. If the cancer is at a stage where it can be removed with surgery, chemotherapy is often used before or after the surgery (see page 29).

When surgery cannot remove the cancer, chemotherapy can be used as a palliative treatment to slow the growth of the cancer and relieve symptoms. For some stage 3 cancers, chemotherapy may be combined with radiation therapy. This is called chemoradiation.

The chemotherapy drugs are usually given as a liquid through a drip inserted into a vein in the arm (intravenous infusion) or as tablets that you swallow. To avoid damaging the veins in your arm, it may also be given through a tube implanted into a vein (called a port, catheter, central line or PICC line). This will stay in place until all your chemotherapy treatment is over.

You will usually receive treatment as an outpatient (not admitted to hospital). Typically, you will have several courses of treatment, with rest periods of a few weeks in between. Your medical oncologist will assess how the treatment is working based on your symptoms and wellbeing, as well as scans and blood tests. Tell your team about any prescription, over-the-counter or natural medicines you are taking, as these may affect how the chemotherapy works in your body.

*“I found chemo a bit daunting – walking into the room with the chairs lined up. But the nurses talked through it with me so I knew what to expect.”* CHERYL

## Side effects of chemotherapy

Chemotherapy can affect healthy cells in the body, which may cause side effects. Some people have few side effects, while others have many. The side effects will depend on the drugs used and the dose. Your medical oncologist and chemotherapy nurses will explain the possible side effects to you, the best ways to manage them and who to contact if you need support.

Side effects of chemotherapy may include feeling very tired (fatigue); feeling sick (nausea); vomiting; mouth ulcers and skin rashes; hair loss; diarrhoea or constipation; flu-like symptoms such as fever, headache and muscle soreness; and poor appetite.

Chemotherapy can also affect the number of cells in your blood. Fewer white blood cells can mean you are more likely to catch infections. Fewer red blood cells (anaemia) can leave you weak and breathless.

Most chemotherapy side effects are temporary and can be managed, so discuss how you are feeling with your treatment team. If you are having chemotherapy as a palliative treatment, your treatment team will help you weigh up the benefits of improving your cancer symptoms against any side effects the chemotherapy is causing.

► See our *Understanding Chemotherapy* booklet.



Many people with pancreatic cancer experience fatigue. This is different to normal tiredness as it doesn't always go away with rest or sleep. The fatigue may be a side effect of treatment or caused by the cancer itself. Managing fatigue is an important part of cancer care. For more information and tips, see our *Understanding Fatigue and Cancer* fact sheet and listen to our "Managing Cancer Fatigue" podcast episode.

## Targeted therapy and immunotherapy

---

Other cancer drug treatments include targeted therapy and immunotherapy.

Targeted therapy targets specific features of cancer cells to stop the cancer growing and spreading, while immunotherapy uses the body's own immune system to fight cancer.

In Australia, as of March 2026, only one targeted therapy drug (olaparib) is approved for use for pancreatic cancer. It is only suitable for a small number of people. This drug has been shown to provide some benefit for people with metastatic pancreatic

cancer who have the BRCA gene changes. Your doctors will be able to provide the latest information about its availability.

So far, no other targeted therapy and immunotherapy drugs have been found to be effective in pancreatic cancer, but research is continuing.

Talk to your doctor about whether a clinical trial is an option for you.

► For more information, see our *Understanding Targeted Therapy* fact sheet.

---

## Radiation therapy

Also known as radiotherapy, radiation therapy uses a controlled dose of radiation to kill cancer cells or injure them so they cannot multiply. The radiation is usually in the form of focused x-ray beams targeted at the cancer. Treatment is painless and carefully planned to do as little harm as possible to healthy body tissue near the cancer.

**Chemoradiation** – For stage 3 cancers that cannot be removed with surgery (locally advanced cancers), radiation therapy may be given with chemotherapy to slow the growth of the cancer. This is called chemoradiation. For cancers that are at a stage where they can be removed by surgery, chemoradiation may also be used before or after the surgery (see page 29).


In chemoradiation, the radiation therapy is delivered over several treatments (called fractions). Each fraction delivers a small dose of radiation that adds up to the total treatment dose. Your doctor will let you know your treatment schedule. Many people have treatment as an outpatient once a day, Monday to Friday, for up to 6 weeks.

Each radiation therapy session takes 10–15 minutes. You will lie on a table under a machine called a linear accelerator that delivers radiation to the affected parts of your body. The machine does not touch you, but it may rotate around you to deliver radiation to the area with cancer from different angles. This allows the radiation to target the cancer more precisely and limits the radiation given to surrounding tissues.

## Linear accelerator (LINAC)

---

This is a general illustration of a LINAC. The machine used for your treatment may look different. A LINAC is large and is often kept in a separate room. An imaging device, such as a CT scan machine, is usually attached to the LINAC. This helps position you accurately on the couch so the radiation targets the correct area of your body.



**Radiation therapy on its own** – Radiation therapy may also be used on its own over shorter periods to relieve symptoms. For example, if a tumour is pressing on a nerve or another organ and causing pain or bleeding, a few doses of radiation therapy may shrink the tumour enough to relieve the symptoms.

**SBRT** – Stereotactic body radiation therapy (SBRT) is an emerging treatment for pancreatic cancer. It delivers a higher dose of radiation per treatment session over fewer treatment sessions than usual radiation therapy. SBRT is not standard practice for pancreatic cancer but may be a treatment option as part of a clinical trial at some cancer centres.

## Side effects of radiation therapy

Radiation therapy can cause side effects, which are mainly related to the area treated. For pancreatic cancer, the treatment is targeted at the abdomen.

Side effects of radiation therapy to the abdomen may include:

- tiredness
- nausea and vomiting
- diarrhoea
- poor appetite
- reflux (when stomach acid flows up into the oesophagus)
- skin irritation.

Most side effects start to improve a few weeks after treatment, but some can last longer or appear later. Late side effects are uncommon, but may include damage to the liver, kidneys, stomach or small bowel. Talk to your radiation oncologist or radiation oncology nurse about ways to manage these side effects.

► See our *Understanding Radiation Therapy* booklet.

## Managing pain in pancreatic cancer

---

If pain becomes an issue, you may need a combination of treatments to achieve good pain control. Options for relieving pain may include:

- strong pain medicines such as opioids
- nerve blocks – when anaesthetic or alcohol is injected into nerves
- anticonvulsant medicines to help control nerve pain
- chemotherapy and/or radiation therapy to shrink cancer pressing on nerves

- complementary therapies such as acupuncture, massage and relaxation techniques.

Tell your treatment team if you have any pain, as it is easier to control when treated early. Don't wait until the pain is severe. Your team can also refer you to a pain specialist or palliative care specialist if needed.

- ▶ See our *Understanding Cancer Pain* booklet.
- 

## How palliative care can help

The treatment options described in this chapter are generally part of palliative treatment (see page 38). Palliative treatment is one aspect of palliative care, in which a team of health professionals work together to meet your physical, emotional, cultural, spiritual and social needs. Specialist palliative care services see people with more complex needs and can also advise other health professionals. See opposite page for more information about palliative care.

Contacting a specialist palliative care service soon after diagnosis gives them the opportunity to get to know you, your family and your circumstances. You can ask your treating doctor for a referral.

- ▶ See our *Understanding Palliative Care* and *Living with Advanced Cancer* booklets and listen to our podcast series *The Thing About Advanced Cancer*.

## How palliative care works

Palliative care is an important part of care for many people with pancreatic cancer. Palliative care may be helpful at any time after diagnosis and is often delivered alongside active cancer treatment to help maintain your quality of life.



### When to start

Palliative care is useful at all stages of advanced cancer. Contacting a palliative care team after a diagnosis of advanced pancreatic cancer can help you work out when is best to start palliative treatment.



### Symptom relief

Palliative treatment can help you manage symptoms related to the cancer or its treatment, such as fatigue, pain or nausea (see pages 42, 46 and 54).



### Who provides care

The palliative care team is made up of people with different skills to help you with a range of issues. Care may be led by a GP, nurse practitioner or community nurse, or a specialist palliative care team.



### Support services

The palliative care team will help you work out how to live in the most fulfilling way you can. They can refer you to organisations and services that can assist with practical, financial and emotional needs.



### Family and carers

If you agree, the palliative care team will involve your family and carers in decisions about your care. They can also provide them with practical and emotional support.

## Key points about treatment to manage symptoms

---

### Treatment goal

The aim of treatment for most locally advanced and advanced pancreatic cancers is to control the cancer for as long as possible, slow down its spread and relieve symptoms. This is known as palliative treatment.

---

### Treatment options

- Advanced pancreatic cancer can be treated palliatively with surgery, chemotherapy or radiation therapy to ease symptoms.
  - Some people have a small tube (stent) inserted to relieve a blockage in the common bile duct or duodenum. This allows bile to flow into the bile duct, or food to move into the small bowel.
  - In some cases, the surgeon will relieve a blockage with an operation to rearrange parts of the digestive system so they avoid the obstruction.
  - Chemotherapy and radiation therapy may be used on their own to relieve symptoms. Sometimes they are combined (chemoradiation) to slow the growth of locally advanced cancer.
  - If you have pain, it can be relieved with pain medicines or palliative cancer treatments. Complementary therapies may also help you cope better with pain and other symptoms.
  - Palliative care helps to improve your quality of life by addressing your physical, emotional, cultural, spiritual and social needs. It also supports families and carers.
-

# Managing your diet and nutrition

Pancreatic cancer and its treatment can affect your ability to eat, digest and absorb essential nutrients. This chapter explains some common diet and nutrition problems, and how to manage them. During and after treatment, it's important to make sure you are eating and drinking enough to maintain your weight and avoid malnutrition or dehydration. Different foods can affect people differently, so you will need to experiment to work out which foods are easiest to digest. Most people will also have to take a special tablet to help with digestion.

▶ See our *Nutrition for People Living with Cancer* booklet for more tips.

## Coping with the changes

Changes to the way you eat may make you feel anxious, particularly when you know eating well is important. Some people find it difficult to cope emotionally with the changes to how and what they can eat. Finding ways to enjoy your meals can help to improve your quality of life. It may help to talk about how you feel with your family and friends. You can also call Cancer Council 13 11 20 – our experienced health professionals may be able to arrange for you to speak with a Cancer Connect volunteer who has had a similar cancer experience.

Some people find that complementary therapies such as relaxation, meditation and acupuncture help them cope with diet and nutrition problems. Always tell your cancer care team if you are using or would like to try any complementary therapies.

▶ See our *Emotions and Cancer* and *Understanding Complementary Therapies* booklets.

## Seeing a dietitian

If you have ongoing problems with food and eating, talk to a dietitian. Dietitians are experts in nutrition who can give you specialist advice on how to cope with nutrition-related problems and eating difficulties throughout different phases of the disease. A dietitian can prepare eating plans for you and give you advice about nutritional supplements.

Dietitians work in all public and most private hospitals. There may be a dietitian connected to your cancer treatment centre – check with your specialist or cancer care coordinator. Dietitians Australia can also help you find an accredited practising dietitian who works in your area and specialises in cancer. For details, call them on 1800 812 942 or visit [dietitiansaustralia.org.au](http://dietitiansaustralia.org.au).

If your GP refers you to a dietitian, you may be eligible for a Medicare rebate to help cover the cost. If you have private health insurance, you may be able to claim part of the cost.

## Nutritional supplements

---

If you can't eat a balanced diet or are losing too much weight, your doctor or dietitian may suggest nutritional supplements such as Sustagen, Ensure, Fortisip or Resource. These provide energy, protein and other nutrients that can help you maintain your strength.

A dietitian can recommend the right nutritional supplement for you and let you know where to buy it

and how much to use. Nutritional supplements should be taken in addition to the foods you are able to eat, and are best used as snacks between meals.

Supplements are available as:

- ready-made drinks, bars, puddings and custards
- powders to mix with milk or water, or to sprinkle on food.

## Enzyme replacement therapy

The pancreas produces digestive enzymes to help break down food. When you have pancreatic cancer, or have had pancreatic surgery, your body may not be able to make enough of these digestive enzymes. This will affect your ability to digest food, particularly fat and protein, and to absorb vital nutrients.

This is often called pancreatic exocrine insufficiency (PEI). Signs of PEI include abdominal pain; bloating and excessive wind; diarrhoea or pale, oily stools that are frothy, loose and difficult to flush; and weight loss.

To help prevent these symptoms, your doctor will prescribe pancreatic enzymes (e.g. Creon). This therapy may be called pancreatic enzyme replacement therapy or PERT. Sometimes, acid-suppressing medicines are also used. The PERT dose will be adjusted depending on your symptoms and diet. It may take time to get this balance right. A dietitian can help you and your doctor work out the correct dose of PERT.




### Taking enzyme supplements

- Take enzyme capsules with water and the first mouthful of food to ensure adequate mixing. With larger meals or meals that take longer to consume (more than 15–20 minutes), you may also need to take them halfway through the meal.
- Always take enzymes when having food or drink that contains fat or protein. Slightly higher doses may be needed with high-fat meals (e.g. fried foods, pizza). You don't need to take enzymes for simple carbohydrates that digest easily (e.g. fruit, fruit juice, black tea, coffee).
- Take enzyme supplements as prescribed. Do not change the dose without first talking to your doctor or dietitian.

## Tips for maintaining your weight


During and after treatment for pancreatic cancer, changes to what you can eat, how you feel about eating and how your body absorbs food can lead to unplanned

### Have regular meals



Eat small meals frequently (e.g. every 2–3 hours), and have a regular eating pattern rather than waiting until you're hungry. Keep ready-to-eat food on hand for when you are too tired to cook (e.g. tinned fruit, yoghurt, frozen meals).

### Choose nourishing food and drink



Ensure that meals and snacks are nourishing and include protein such as meat, chicken, fish, dairy products, legumes (e.g. lentils, chickpeas), eggs, tofu, nuts and nut butters. Choose nourishing drinks (e.g. milk, smoothies). A dietitian may also suggest nutritional supplement drinks (see page 50).

### Load up your food



There are different ways to add extra kilojoules to your food:

- Add milk powder to cereals, sauces, mashed vegetables, soup, drinks, egg dishes and desserts.
- Add cheese to sauces, soup, baked beans, vegetables, casseroles, salads and egg dishes.
- Add sugar, honey or golden syrup to cereals, porridge or drinks.
- Use a powdered nutritional supplement recommended by your dietitian (see page 50).


weight loss. This can cause a loss of strength, increase fatigue and affect how you cope with treatment. The tips below and on the following pages may help.

### Adjust to taste changes



You may find your sense of taste changes after treatment. If food tastes bland, add extra flavouring such as herbs, lemon, lime, ginger, garlic, honey, chilli, pepper, Worcestershire sauce, soy sauce or pickles. Eating moist fruits such as berries or sucking boiled lollies can help if you have a bitter or metallic taste in your mouth.

### Avoid strong food smells



If food smells bother you, ask family or friends to do the cooking. You may also prefer cold food or food at room temperature without a strong smell.

### Talk to a dietitian



A dietitian can help if you are finding it hard to work out the right foods to help gain weight. You should also check with a dietitian before cutting out particular foods.

### Follow your appetite



It's okay to focus on eating foods that you enjoy. Gaining or maintaining weight is more important at the moment than avoiding extra fat and sugar.

## Nausea and vomiting

Nausea and vomiting can occur because of the cancer or its treatment. For some people, just the thought of treatment or eating or the smell of food can make them feel unwell. There is a range of anti-nausea medicines (antiemetics) that you can take regularly to control symptoms. If the one you are prescribed doesn't work, let your doctor or nurse know so you can try another medicine.

Also advise your treatment team if vomiting lasts for more than a day or if you can't keep any fluids down, as you may become dehydrated. Signs of dehydration include a dry mouth, dark urine, dizziness and confusion.

If you have persistent vomiting, the duodenum (the first part of the small bowel) may be blocked, so see your doctor as soon as possible. You may need surgery to clear the blockage (see pages 39–40).



### Coping with nausea

- Eat and drink slowly. Chew food well.
- Some anti-nausea medicines need to be taken half an hour before meals – ask your doctor about this.
- Try to eat a little bit at regular intervals – not eating or skipping meals can make nausea worse.
- Suck on boiled lollies. Try peppermint-flavoured or lemon-flavoured lollies.
- Avoid strong odours and cooking smells.
- Snack on bland foods such as dry crackers or toast.
- Drink ginger beer, ginger ale or ginger tea, or suck on candied ginger.
- See opposite page for tips on dealing with vomiting.
- ▶ Listen to our “Appetite Loss and Nausea” podcast episode at [cancercouncil.com.au/podcasts](http://cancercouncil.com.au/podcasts).

## Steps to recovery after vomiting



### 1. Take small sips

Don't try to force down food. Sip small amounts of liquid as often as possible. Try flat dry ginger ale, cold flat lemonade, weak cordial, or cold apple or orange juice.




### 2. Introduce nourishing fluids

If the vomiting has stopped but you still feel sick, slowly introduce more nourishing fluids. Start with cold or iced drinks. Prepare milk or fruit drinks with some water so they are not too strong. You can also try diluted fluids such as clear broth or weak tea.



### 3. Start solid food

Next, try to eat small amounts of solid foods, such as plain dry biscuits, toast or bread with honey or jam, or smooth congee (rice porridge). Stewed fruits and yoghurt are also good. Aim to eat small amounts of food often, rather than 3 large meals a day.



### 4. Return to a normal diet

As soon as you can, increase your food intake until you are eating a normal, balanced diet. Limit rich foods, such as fatty meats or full-cream dairy products. Your doctor or dietitian may suggest adding extra nourishment (such as nutritional supplements) on your good days to make up for the days you can't eat properly.

# Diarrhoea

Diarrhoea is when your bowel motions become loose, watery and frequent. You may also get abdominal cramping, wind and pain.

Pancreatic cancer treatments, other medicines, infections, reactions to certain foods, and anxiety can all cause diarrhoea.

If the tips listed below don't work for you, talk to your doctor about trying anti-diarrhoea medicine.

You should also let your doctor know if your stools are pale in colour, oily, very smelly, float and are difficult to flush, or you notice an oily film floating in the toilet. This may be a sign that you do not have enough pancreatic enzymes. You may need to start pancreatic enzyme replacement therapy (PERT) or adjust your dose. See page 51 for more information.



## Managing diarrhoea

- Drink plenty of liquids (e.g. water, fruit juice, weak cordial) to replace lost fluids.
- Try lactose-reduced milk or soy milk if you develop a temporary intolerance to the sugar (lactose) in milk. This can sometimes occur when you have diarrhoea. Small amounts of hard cheese and yoghurt are usually okay.
- Avoid alcohol and limit caffeine and spicy foods as these can all make diarrhoea worse.
- Always let your treatment team know if you notice any changes to your bowel habits.

# Diabetes

Insulin is a hormone that controls the amount of sugar in the blood. Diabetes, or high blood sugar levels, can occur if your pancreas is not making enough insulin. This is why some people develop diabetes shortly before pancreatic cancer is diagnosed (when the cancer is affecting how much insulin the pancreas can make) or soon after surgery (when some or all of the pancreas has been removed).

The way diabetes is managed varies from person to person but often includes both dietary changes and insulin injections. Sometimes medicines are given as tablets that you swallow.

Your GP can help you manage the condition, but you may be referred to an endocrinologist, a specialist in hormone disorders. You may also be referred to a dietitian for help with your diet and to a diabetes specialist nurse, who can help to coordinate your care and provide support.



## Coping with diabetes

- Eat small meals and snacks regularly to help control blood sugar levels.
- Talk to your endocrinologist or GP about medicines to help control the diabetes.
- If you are taking diabetes medicine, include high-fibre carbohydrate foods at every meal to avoid low blood sugar levels. Wholegrain breads and cereals, vegetables and fruit are all suitable foods.
- Talk to your doctors and dietitian for more information about diabetes.
- Get in touch with the National Diabetes Services Scheme on 1800 637 700 or visit [ndss.com.au](http://ndss.com.au). They can provide advice on managing diabetes.

# Looking after yourself

Cancer can cause physical and emotional strain, so it's important to look after your wellbeing. Cancer Council has free booklets and programs to help you during and after treatment. Call 13 11 20 to find out more, or visit your local Cancer Council website (see back cover).

**Eating well** – Healthy food can help you cope with treatment and side effects. A dietitian can explain how to manage any special dietary needs or eating problems and choose the best foods for your situation.

▶ See our *Nutrition for People Living with Cancer* booklet.

**Staying active** – Physical activity can reduce tiredness, improve circulation and lift mood. The right exercise for you depends on what you are used to, how you feel, and your doctor's advice.

▶ See our *Exercise for People Living with Cancer* booklet.

**Complementary therapies** – Complementary therapies are designed to be used alongside conventional medical treatments. Therapies such as massage, relaxation and acupuncture can increase your sense of control, decrease stress and anxiety, and improve your mood. Let your doctor know about any therapies you are using or thinking about trying, as some may not be safe or evidence-based.

▶ See our *Understanding Complementary Therapies* booklet.



Alternative therapies are therapies used instead of conventional medical treatments. These are unlikely to be scientifically tested, may prevent successful treatment of the cancer and can be harmful. Cancer Council does not recommend the use of alternative therapies as a cancer treatment.

**Work and money** – Cancer can change your financial situation, especially if you have extra medical expenses or need to stop working. Getting professional financial advice and talking to your employer can give you peace of mind. You can also check whether any financial assistance is available to you by asking a social worker at your hospital or treatment centre or calling Cancer Council 13 11 20.

▶ See our *Cancer and Your Finances* and *Cancer, Work and You* booklets.

**Relationships** – Having cancer can affect your relationships with family, friends and colleagues in different ways. Cancer is stressful, tiring and upsetting, and this may strain relationships. The experience of cancer may also result in positive changes to your values, priorities or outlook on life. Give yourself time to adjust to what's happening, and do the same for those around you. It may help to discuss your feelings with each other.

▶ See our *Emotions and Cancer* booklet.

**Sex life** – Cancer can affect your sex life in physical and emotional ways. The impact of these changes depends on many factors, such as treatment and side effects, your self-confidence, and if you have a partner. Although sexual intercourse may not always be possible, closeness and sharing can still be part of your relationship.

▶ See our *Sex, Intimacy and Cancer* booklet.

**Contraception and fertility** – If you can have sex, you may need to use certain types of contraception to protect your partner or avoid pregnancy for a time. Your doctor will explain what precautions to take. They will also tell you if treatment will affect your fertility permanently or temporarily. If having children is important to you, discuss the options with your doctor before starting treatment.

▶ See our *Fertility and Cancer* booklet.

# Living with pancreatic cancer

Life after a diagnosis of pancreatic cancer can present many challenges. It is important to allow yourself time to adjust to the physical and emotional changes. Establish a new daily routine that suits you and the symptoms you're coping with. Your family and friends may also need time to adjust.

For some people, the cancer goes away with treatment. Other people will have ongoing treatment to manage symptoms. You are likely to feel a range of emotions about having pancreatic cancer. Talk to your treatment team if you are finding it hard to manage your emotions. Cancer Council 13 11 20 can also provide you with some strategies for coping with the emotional and practical aspects of living with pancreatic cancer.

► See our *Emotions and Cancer* booklet.

## Dealing with feelings of sadness

---

If you have continued feelings of sadness, have trouble getting up in the morning or have lost motivation to do things that previously gave you pleasure, you may be experiencing depression. This is quite common among people who have cancer.

Talk to your GP, because counselling or medication – even for a short time – may help. Some people can

get a Medicare rebate for sessions with a psychologist. Cancer Council may also run a counselling program in your area.

For information about coping with depression and anxiety, call Beyond Blue on 1300 22 4636 or visit [beyondblue.org.au](http://beyondblue.org.au). For 24-hour crisis support, call Lifeline 13 11 14 or visit [lifeline.org.au](http://lifeline.org.au).



For information and insights that can help you navigate through the challenges of living with advanced cancer, listen to Cancer Council's podcast series *The Thing About Advanced Cancer*. To listen, go to [cancercouncil.com.au/podcasts](http://cancercouncil.com.au/podcasts).

## When the cancer is advanced

If you are told that you have advanced pancreatic cancer, it may bring up different emotions and reactions. You may not know what to say or think; you may feel sadness, anger, disbelief or fear.

There is no right or wrong way to react. Give yourself time to take in what is happening and accept that some days will be easier than others.

Many people diagnosed with pancreatic cancer think about what will happen to them if or when the disease progresses. You may question how much more time you have to live and begin going over your life and what it has meant for you. These thoughts are natural in this situation.

You might find it helpful to talk to your GP and the palliative care team about what you are going through.

In addition to the palliative care doctors and nurses, the specialist palliative care team may include a social worker, counsellor or spiritual care practitioner (pastoral carer). They can explain what to expect and how any symptoms will be managed.

If you are not already in contact with a palliative care service, talk to your cancer specialist about a referral.

- ▶ See our *Living with Advanced Cancer* and *Understanding Palliative Care* booklets.

# Seeking support

A cancer diagnosis can affect every aspect of your life. You will probably experience a range of emotions – fear, sadness, anxiety, anger and frustration are all common reactions. Cancer also often creates practical and financial issues.

There are many sources of support and information to help you, your family and carers navigate all stages of the cancer experience, including:

- information about cancer and its treatment
- access to benefits and programs to ease the financial impact of cancer treatment
- home care services, such as Meals on Wheels, visiting nurses and home help
- aids and appliances
- support groups and programs
- counselling services.

The availability of services may vary depending on where you live, and some services will be free but others might have a cost.

To find good sources of support and information, you can talk to the social worker or nurse at your hospital or treatment centre, or get in touch with Cancer Council 13 11 20.

*“My family members don’t really understand what it’s like to have cancer thrown at you, but in my support group, I don’t feel like I have to explain.”* SAM

## Support from Cancer Council

Cancer Council offers a range of services to support people affected by cancer, their families and friends. Services may vary by location.

### Cancer Council 13 11 20



Our experienced health professionals will answer any questions you have about your situation and link you to local services (see inside back cover).

### Information resources



Cancer Council produces booklets and fact sheets on more than 40 types of cancer, as well as treatments, emotional and practical issues, and recovery. Call 13 11 20 or visit your local Cancer Council website.

### Legal and financial support



If you need advice on legal or financial issues, we may be able to refer you to qualified professionals. These services are free for people who can't afford to pay. Financial assistance may also be available. To find out more, call Cancer Council 13 11 20.

### Practical help



Cancer Council can help you find services or offer guidance to manage the practical impacts of cancer. This may include helping you access accommodation and transport services.

### Peer support services



You might find it helpful to share your thoughts and experiences with other people affected by cancer. Cancer Council can link you with individuals or support groups by phone, in person, or online. Call 13 11 20 or visit [cancercouncil.com.au/OC](http://cancercouncil.com.au/OC).

## Useful websites

You can find many useful resources online, but not all websites are reliable. These websites are good sources of support and information.

### Australian

Cancer Council Australia	<a href="https://cancer.org.au">cancer.org.au</a>
Cancer Council Online Community	<a href="https://cancercouncil.com.au/OC">cancercouncil.com.au/OC</a>
Cancer Council Podcasts	<a href="https://cancercouncil.com.au/podcasts">cancercouncil.com.au/podcasts</a>
Guides to Best Cancer Care	<a href="https://cancer.org.au/cancercareguides">cancer.org.au/cancercareguides</a>
Australian Pancreatic Cancer Genome Initiative	<a href="https://pancreaticcancer.net.au">pancreaticcancer.net.au</a>
Cancer Australia	<a href="https://canceraustralia.gov.au">canceraustralia.gov.au</a>
Carer Gateway	<a href="https://carergateway.gov.au">carergateway.gov.au</a>
Dietitians Australia	<a href="https://dietitiansaustralia.org.au">dietitiansaustralia.org.au</a>
eviQ (cancer treatment information)	<a href="https://eviq.org.au">eviq.org.au</a>
GI Cancer Trials	<a href="https://gicancer.org.au">gicancer.org.au</a>
Palliative Care Australia	<a href="https://palliativecare.org.au">palliativecare.org.au</a>
Pancare Foundation	<a href="https://pancare.org.au">pancare.org.au</a>
PanKind	<a href="https://pankind.org.au">pankind.org.au</a>
Pancare Support	<a href="https://pancare.org.au/support">pancare.org.au/support</a>
Radiation Oncology: Targeting Cancer	<a href="https://targetingcancer.com.au">targetingcancer.com.au</a>
Services Australia	<a href="https://servicesaustralia.gov.au">servicesaustralia.gov.au</a>

### International

American Cancer Society	<a href="https://cancer.org">cancer.org</a>
Cancer Research UK	<a href="https://cancerresearchuk.org">cancerresearchuk.org</a>
Macmillan Cancer Support (UK)	<a href="https://macmillan.org.uk">macmillan.org.uk</a>

# Caring for someone with cancer

You may be reading this booklet because you are caring for someone with cancer. What this means for you will vary depending on the situation. Being a carer can bring a sense of satisfaction, but it can also be challenging and stressful.

It is important to look after your own physical and emotional wellbeing. Give yourself some time out and share your concerns with somebody neutral such as a counsellor or your doctor, or try calling Cancer Council 13 11 20. There is a wide range of support available to help you with the practical and emotional aspects of your caring role.

**Support services** – Support services such as Meals on Wheels, home help or visiting nurses can help you in your caring role. You can find local services, as well as information and resources, through the Carer Gateway. Call 1800 422 737 or visit [carergateway.gov.au](http://carergateway.gov.au).

**Support groups and programs** – Many cancer support groups and cancer education programs are open to carers as well as to people with cancer. Support groups and programs offer the chance to share experiences and ways of coping.

**Carers Australia** – Carers Australia provides information and advocacy for carers. Visit [carersaustralia.com.au](http://carersaustralia.com.au).

**Cancer Council** – You can call Cancer Council 13 11 20 or visit your local Cancer Council website to find out more about carers' services.

▶ See our *Caring for Someone with Cancer* booklet.

## Question checklist

---

Asking your doctor questions will help you make an informed choice. You may want to include some of the questions below in your own list.

---

### Diagnosis

- What type of pancreatic cancer do I have?
  - Has the cancer spread? If so, where has it spread? How fast is it growing?
  - Are the latest tests and treatments for this cancer available in this hospital?
  - Will a multidisciplinary team be involved in my care?
  - Are there clinical guidelines for this type of cancer?
- 

### Treatment

- What treatment do you recommend? What is the aim of the treatment?
  - Are there other treatment choices for me? If not, why not?
  - If I don't have the treatment, what should I expect?
  - How long do I have to make a decision?
  - I'm thinking of getting a second opinion. Can you recommend anyone?
  - How long will treatment take? Will I have to stay in hospital?
  - Are there any out-of-pocket expenses not covered by Medicare or my private health cover? Can the cost be reduced if I can't afford it?
  - How will we know if the treatment is working?
  - Are there any clinical trials or research studies I could join?
  - Do I need the support of a specialist palliative care service?
- 

### Side effects

- What are the risks and possible side effects of each treatment?
  - Will I have a lot of pain? What will be done about this?
  - Can I work, drive and do my normal activities while having treatment?
  - Will the treatment affect my sex life and fertility?
  - Should I change my diet or physical activity during or after treatment?
  - Are there any complementary therapies that might help me?
- 

### After treatment for early pancreatic cancer

- How often will I need check-ups after treatment?
  - If the cancer returns, how will I know? What treatments could I have?
-

# Glossary

---

## **abdomen**

The part of the body between the chest and hips, which contains the stomach, spleen, pancreas, liver, gallbladder, bowel, bladder and kidneys. Also known as the belly.

## **adenocarcinoma**

Cancer that starts in the mucus-producing (glandular) cells that form part of the lining of internal organs. Pancreatic adenocarcinoma is the most common pancreatic cancer. It starts in the pancreatic duct lining.

## **adjuvant therapy**

A treatment given after the main treatment to lower the risk that the cancer will come back.

## **advanced cancer**

Cancer that is unlikely to be cured. In most cases, the cancer has spread to other parts of the body (secondary or metastatic cancer). Treatment can often still control the cancer and manage symptoms.

## **anaesthetic**

A drug that stops a person feeling pain during a medical procedure. Local and regional anaesthetics numb part of the body; a general anaesthetic causes a temporary loss of consciousness.

---

## **benign**

Not cancerous or malignant.

## **bile**

A substance produced by the liver and stored in the gallbladder. Bile helps the digestive system break down fats from food.

## **bilirubin**

A dark yellow-brown substance found in bile. If bilirubin builds up in the body, it can cause symptoms of jaundice. See also jaundice.

## **biopsy**

The removal of a sample of cells or tissue from the body for examination under a microscope to help diagnose a disease.

## **borderline resectable cancer**

Cancer that could be removed with surgery if other treatment can shrink the cancer first.

---

## **catheter**

A hollow, flexible tube through which fluids can be passed into the body or drained from it.

## **cells**

The basic building blocks of the body. A human is made of billions of cells adapted for different functions.

## **chemoradiation**

Treatment that combines chemotherapy with radiation therapy. This may also be known as chemoradiotherapy.

## **chemotherapy**

A cancer treatment that uses drugs to kill cancer cells or slow their growth. May be given alone or in combination with other treatments.

## **cirrhosis**

A condition in which healthy liver cells are replaced by scar tissue.

## **clinical trial**

A research study that tests new approaches to prevention, screening, diagnosis or treatment, to see if they are better than current approaches.

## **common bile duct**

The tube through which bile travels from the liver and gallbladder to the bowel.

## **constipation**

Difficulty passing a bowel motion regularly or often.

### **CT scan**

Computerised tomography scan. This scan uses x-rays to create a detailed cross-sectional picture of the body.

---

### **diabetes**

A condition in which sugars are not taken up in the body properly because the pancreas does not make enough of the necessary hormone (insulin), or the body has become resistant to the effect of insulin. In pancreatic cancer, diabetes sometimes occurs shortly before a diagnosis or as a result of treatment.

### **diarrhoea**

Opening the bowels very frequently. Motions may be loose and watery.

### **dietitian**

A university-qualified health professional who supports and educates patients about nutrition and diet during cancer treatment and recovery.

### **distal pancreatectomy**

Surgery to remove tumours in the tail and body of the pancreas.

### **duct**

An enclosed tube or passage in the body.

### **duodenum**

The first section of the small bowel.

---

### **endocrine gland**

A gland that releases hormones that control the amount of sugar in the blood.

### **endocrine system**

The system of the body that is made of glands that produce hormones.

### **endocrinologist**

A doctor who specialises in treating people with disorders of the endocrine system.

### **endoscope**

A flexible tube with a light and camera on the end. It is used for a diagnostic test known as an endoscopy (see next column).

### **endoscopic retrograde cholangiopancreatography (ERCP)**

A procedure using an endoscope to check the pancreas and bile ducts, or to insert a stent.

### **endoscopic ultrasound (EUS)**

A test using an endoscope with an ultrasound probe on the end. The probe releases soundwaves that are translated into a picture on a computer screen.

### **endoscopy**

A type of internal examination or diagnostic test using an endoscope (see previous column) to examine the inside of the body.

### **enzymes**

Proteins that aid digestion and are essential for the normal functioning of the body.

### **exocrine tumour**

A tumour that starts in the cells that make pancreatic enzymes to aid digestion. More than 95% of pancreatic tumours are exocrine tumours, and most of these are adenocarcinomas.

---

### **fatigue**

Extreme feeling of tiredness and lack of energy that may not go away with rest.

---

### **gallbladder**

A small, pear-shaped organ on the underside of the liver that stores bile.

### **gastroenterologist**

A doctor who specialises in diagnosing and treating disorders of the digestive system.

### **genes**

The microscopic units that determine how the body's cells grow and behave. Genes are found in every cell of the body and are inherited from both parents.

### **genetic/genomic testing**

Tests that look for changes in genes to find out what has caused the cancer and how it might grow or spread. The results can help

doctors work out the best treatment. Genetic tests check for specific genes; genomic tests look at many genes at once. Gene changes can be inherited from family or happen during a person's lifetime (acquired).

### **glands**

Specialised organs or groups of cells that make various fluids that are used in the body or excreted.

### **glucagon**

A hormone that increases blood sugar (glucose) levels.

### **grade**

A score that describes how quickly a tumour is likely to grow.

---

### ***Helicobacter pylori***

Bacteria that can live in the stomach and small bowel and may lead to stomach ulcers and cancer. Also called *H. pylori*.

### **hormones**

Chemicals in the body that send information between cells. Hormones control many of the body's functions, including how you grow, develop and reproduce.

---

### **immunotherapy**

Drugs that use the body's own immune system to fight cancer.

### **insulin**

A hormone released by the pancreas to regulate the amount of sugar (glucose) in the blood. If your body does not produce enough insulin, you will develop diabetes.

### **interventional radiologist**

A specialist doctor who uses imaging scans to diagnose cancer, may perform a biopsy and may deliver some treatments.

---

### **jaundice**

A condition caused by high levels of a substance called bilirubin in the blood. It can

occur when the bile ducts are blocked or the liver is not working properly. Jaundice causes yellow, itchy skin; the whites of the eyes to turn yellow; pale stools (poo); and dark urine.

---

### **keyhole surgery (laparoscopy)**

Surgery done through small cuts in the abdomen using a thin viewing instrument (called a laparoscope) with a light and camera.

---

### **laparoscopy (laparoscopic surgery)**

See keyhole surgery.

### **liver**

A large organ in the top right side of the abdomen. Its functions include making bile, which helps digestion.

### **locally advanced cancer**

Cancer that has spread to nearby organs or blood vessels and often cannot be removed with surgery.

### **lymphatic system**

A network of vessels, nodes and organs that removes excess fluid from tissues, absorbs fatty acids, transports fat and produces immune cells.

### **lymph nodes (lymph glands)**

Small lumps of tissue that collect and destroy bacteria and viruses. They help protect the body against disease and infection.

---

### **magnetic resonance**

### **cholangiopancreatography (MRCP) scan**

A more detailed MRI scan that can check the bile duct for blockages.

### **malignant**

Cancerous. Malignant cells can spread (metastasise) and eventually cause death if they cannot be treated.

### **metastasis (plural: metastases)**

Cancer that has spread from a primary cancer in another part of the body. Also known as secondary or advanced cancer.

### **MRI scan**

Magnetic resonance imaging scan. These scans use magnetic fields and radio waves to take detailed, cross-sectional pictures of the body.

---

### **nausea**

Feeling sick or wanting to be sick.

### **neoadjuvant therapy**

A treatment given before another treatment to make that treatment more successful.

### **neuroendocrine tumour (NET)**

A rare type of tumour that affects the endocrine and nervous systems. Some types of NETs affect the pancreas.

---

### **outpatient**

A person who visits a hospital for medical treatment without being admitted into hospital.

---

### **palliative care**

The holistic care of people who have a life-limiting illness, their families and carers. It aims to maintain quality of life by addressing physical, emotional, cultural, spiritual and social needs. Also known as supportive care.

### **palliative treatment**

Medical treatment for people with advanced cancer to help them manage pain and other physical and emotional symptoms.

### **pancreas**

An organ in the digestive and endocrine systems. The pancreas produces insulin and some of the enzymes needed to digest food.

### **pancreatectomy**

The removal of all or part of the pancreas.

### **pancreatic duct**

A tube that connects the pancreas to the first part of the small bowel (duodenum).

### **pancreatic enzyme replacement therapy (PERT)**

This replaces the enzymes that your pancreas would normally make. It helps to digest your food by breaking down carbohydrates, fats and proteins in your food. The enzymes come in capsules that you swallow. PERT is also known by the brand name Creon.

### **pancreatic exocrine insufficiency (PEI)**

When the body does not make enough digestive enzymes to properly digest food.

### **pancreatic neuroendocrine tumour (pancreatic NET or PNET)**

A neuroendocrine tumour affecting the pancreas.

### **pancreaticoduodenectomy**

See Whipple procedure.

### **pancreatitis**

Inflammation of the pancreas.

### **peritoneum**

The lining of the abdomen.

### **PET-CT scan**

A positron emission tomography (PET) scan combined with a CT scan. In a PET scan, the person is injected with a small amount of radioactive solution. This makes cancerous areas show up brighter on the scan.

### **prognosis**

The expected outcome of a particular person's disease.

---

### **radiation therapy**

The use of targeted radiation to kill or damage cancer cells so they cannot grow, multiply or spread.

### **recurrence**

The return of a disease after a period of improvement (remission).

### **resectable cancer**

Cancer that can be removed with surgery.

---

### stage

The extent of a cancer and whether it has spread from an original site to other parts of the body.

### stent

A metal or plastic tube placed into a blocked organ to create a passage for substances to pass through.

### stools (faeces or poo)

The bulky mass of waste material that normally leaves the body through the anus.

---

### targeted therapy

Drugs that target specific features of cancer cells to stop cancer growing and spreading.

### tissue

A collection of cells of similar type that make up an organ or structure in the body.

### total pancreatectomy

The removal of the entire pancreas. The spleen, gall bladder, bile duct, part of the stomach and small bowel, and nearby lymph nodes are often removed at the same time.

### tumour

A new or abnormal growth of tissue on or in the body; may be benign or malignant.

### tumour marker

A chemical produced by cancer cells and

released into the blood. It may suggest the presence of a tumour in the body. Markers can be found by blood tests or by testing tumour samples.

---

### unresectable

Not able to be removed with surgery.

---

### Whipple procedure

Surgery to remove the head of the pancreas and surrounding structures. Also called a Whipple operation, Whipple surgery or pancreaticoduodenectomy.

---

### Can't find a word here?

For more cancer-related words, visit:

- [cancercouncil.com.au/words](https://cancercouncil.com.au/words)
  - [cancervic.org.au/glossary](https://cancervic.org.au/glossary)
- 

## References

1. RC Gandy et al., "Refining the care of patients with pancreatic cancer: The AGITG Pancreatic Cancer Workshop consensus", *Medical Journal of Australia*, vol. 204, iss. 11, 2016, pp. 419–22.
2. T Conroy et al. on behalf of the ESMO Guidelines Committee, "Pancreatic cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up", *Annals of Oncology*, vol. 34, no. 11, 2023, pp. 987–1002.
3. Australian Institute of Health and Welfare (AIHW), *Cancer Data in Australia*, viewed 24 September 2025, available from [aihw.gov.au/reports/cancer/cancer-data-in-Australia](https://aihw.gov.au/reports/cancer/cancer-data-in-Australia).



# How you can help

At Cancer Council, we're dedicated to improving cancer control. As well as funding millions of dollars in cancer research every year, we advocate for the highest quality care for cancer patients and their families. We create cancer-smart communities by educating people about cancer, its prevention and early detection. We offer a range of practical and support services for people and families affected by cancer. All these programs would not be possible without community support, great and small.

**Join a Cancer Council event:** Join one of our community fundraising events such as Daffodil Day, Australia's Biggest Morning Tea, Relay For Life, Girls' Night In and other Pink events, or hold your own fundraiser or become a volunteer.

**Make a donation:** Any gift, large or small, makes a meaningful contribution to our work in supporting people with cancer and their families now and in the future.

**Buy Cancer Council sun protection products:** Every purchase helps you prevent cancer and contribute financially to our goals.

**Help us speak out for a cancer-smart community:** We are a leading advocate for cancer prevention and improved patient services. You can help us speak out on important cancer issues and help us improve cancer awareness by living and promoting a cancer-smart lifestyle.

**Join a research study:** Cancer Council funds and carries out research investigating the causes, management, outcomes and impacts of different cancers. You may be able to join a study.

To find out more about how you, your family and friends can help, please call your local Cancer Council.



# Cancer Council

## 13 11 20

Being diagnosed with cancer can be overwhelming. At Cancer Council, we understand it isn't just about the treatment or prognosis. Having cancer affects the way you live, work and think. It can also affect our most important relationships.

When disruption and change happen in our lives, talking to someone who understands can make a big difference. Cancer Council has been providing information and support to people affected by cancer for over 50 years.

Calling 13 11 20 gives you access to trustworthy information that is relevant to you. Our experienced health professionals are available to answer your questions and link you to services in your area, such as transport, accommodation and home help. We can also help with other matters, such as legal and financial advice.

If you are finding it hard to navigate through the health care system, or just need someone to listen to your immediate concerns, call 13 11 20 and find out how we can support you, your family and friends.



If you need information in a language other than English, an interpreting service is available. Call 131 450.



If you are deaf, or have a hearing or speech impairment, you can contact us through the National Relay Service. [accesshub.gov.au](http://accesshub.gov.au)

---

*Cancer Council services and programs vary in each area.  
13 11 20 is charged at a local call rate throughout Australia (except from mobiles).*

For information & support  
on cancer-related issues,  
call **Cancer Council 13 11 20**

## Visit your local Cancer Council website

---

**Cancer Council ACT**  
[actcancer.org](http://actcancer.org)

**Cancer Council Queensland**  
[cancerqld.org.au](http://cancerqld.org.au)

**Cancer Council Victoria**  
[cancervic.org.au](http://cancervic.org.au)

**Cancer Council NSW**  
[cancercouncil.com.au](http://cancercouncil.com.au)

**Cancer Council SA**  
[cancersa.org.au](http://cancersa.org.au)

**Cancer Council WA**  
[cancerwa.asn.au](http://cancerwa.asn.au)

**Cancer Council NT**  
[cancer.org.au/nt](http://cancer.org.au/nt)

**Cancer Council Tasmania**  
[cancer.org.au/tas](http://cancer.org.au/tas)

**Cancer Council Australia**  
[cancer.org.au](http://cancer.org.au)

*This booklet is funded through the generosity of the people of Australia.  
To support Cancer Council, call your local Cancer Council or visit your local website.*

