



Submission from Cancer Council Australia to FSANZ's Application A1090- Voluntary addition of vitamin D to breakfast cereal

27 February 2015

Cancer Council is Australia's peak national non-government cancer control organisation. Its members are the eight state and territory cancer organisations, working together to undertake and fund cancer research, prevent and control cancer, and provide information and support for people affected by cancer. Cancer Council's goal is to lead the development and promotion of national cancer control policy in Australia, in order to prevent cancer and reduce the illness, disability and death caused by cancer.

Cancer Council has concerns regarding FSANZ's proposal to permit the addition of vitamin D₂ to breakfast cereal. We note the FSANZ call for submissions on Application A1090 states that Standard 1.1.1 of the Food Standards Code permits two forms of vitamin D: D₂ and D₃ to be added to relevant foods, as they can be considered equivalent at dietary intakes up to 25 µg/day. Therefore FSANZ has proposed to permit vitamin D₂ as well as vitamin D₃ for addition to breakfast cereal.

Recent studies suggest that circulating vitamin D₂ is associated with increased risk of all-cause mortality and inversely associated with vitamin D₃ concentrations. Our organisation does not support the draft variation proposed as it does not differentiate between vitamin D₂ and vitamin D₃ for food fortification.

There are two lines of evidence that indicate vitamin D₂ should not be used as a dietary supplement. Firstly there is evidence from a number of systematic reviews and meta-analyses covering a substantial body of literature, that vitamin D₂ does not raise circulating 25 hydroxy (OH) vitamin D levels to the same extent as does D₃^{1, 2, 3}. Other evidence suggests that D₂ may reduce the availability of D₃ and does not increase calcitriol (1, 25 dihydroxy vitamin D) levels⁴. A recent randomised controlled trial in New Zealand evaluated the effect of 25 µg of D₂ or D₃ against placebo in healthy adults over 25 weeks of supplementation beginning at the end of summer⁵. Although D₂ supplementation maintained a higher serum 25(OH)D than in the placebo group, the fall in 25(OH)D₃ was larger in the D₂ supplementation group than the placebo group. This trial was different from most others as supplementation was started when the circulating 25(OH)D levels were relatively high. The results still clearly indicate the superiority of D₃ compared with D₂ in maintaining vitamin D levels.

Secondly, and of more concern are the meta-analysis by Chowdhury et al⁶ indicating that while vitamin D₃ supplementation decreased all-cause mortality, vitamin D₂ supplementation did not, and in some studies increased mortality risk; and the Cochrane review of vitamin D supplementation and mortality⁷ which found that vitamin D supplementation decreased

mortality, but when vitamin D₂ and vitamin D₃ were considered separately the benefit was only seen with vitamin D₃, and vitamin D₂ was associated with increased risk in trials at high risk of bias or in trials including participants with vitamin D insufficiency.

Researchers from Cancer Council VIC have unpublished data from the Melbourne Collaborative Cohort Study that show circulating vitamin D₂ is associated with increased risk of all-cause mortality and inversely associated with vitamin D₃ concentrations. These data are expected to be published soon and are available to share confidentially with FSANZ before journal publication.

In view of the above data suggesting that vitamin D₂ supplementation would not achieve the expected increase in circulating vitamin D levels, and may actually have a harmful effect there is no justification for allowing vitamin D₂ to be added to any foods. Cancer Council does not support the draft variation proposed by FSANZ - Food Standards (Application A1090 – Addition of Vitamin D to Breakfast Cereal) Variation. Cancer Council also strongly recommends a review of Standard 1.1.1 - Preliminary Provisions which permits two forms of vitamin D: D₂ and D₃ to be added to relevant foods. Given the potential harmful effects of Vitamin D₂, the Food Standards Code needs to differentiate between D₂ and D₃ for food fortification.

While Cancer Council supports the voluntary addition of vitamin D₃ to breakfast cereal, Cancer Council cannot support the current proposal as it permits the addition of vitamin D₂.

For further information on this submission:

Mr Paul Grogan
Director, Public Policy and Advocacy
Cancer Council Australia
paul.grogan@cancer.org.au
T: +61 2 8063 4155

References

1. Houghton LA and Vieth R. The case against ergocalciferol (vitamin D2) as a vitamin supplement. *Am J Clin Nutr.* 2006; 84: 694-7.
2. Tripkovic L, Lambert H, Hart K, Smith CP, Bucca G, Penson S, Chope G, Hypponen E, Berry J, Vieth R and Lanham-New S. Comparison of vitamin D2 and vitamin D3 supplementation in raising serum 25-hydroxyvitamin D status: a systematic review and meta-analysis. *Am J Clin Nutr.* 2012; 95: 1357-64.
3. Autier P, Gandini S and Mullie P. A systematic review: influence of vitamin D supplementation on serum 25-hydroxyvitamin D concentration. *J Clin Endocrinol Metab.* 2012; 97: 2606-13.
4. Swanson CM, Nielson CM, Shrestha S, Lee CG, Barrett-Connor E, Jans I, Cauley JA, Boonen S, Bouillon R, Vanderschueren D and Orwoll ES. Higher 25(OH)D2 is associated with lower 25(OH)D3 and 1,25(OH)2D3. *J Clin Endocrinol Metab.* 2014; 99: 2736-44.
5. Logan VF, Gray AR, Peddie MC, Harper MJ and Houghton LA. Long-term vitamin D3 supplementation is more effective than vitamin D2 in maintaining serum 25-hydroxyvitamin D status over the winter months. *Br J Nutr.* 2013; 109: 1082-8.
6. Chowdhury R, Kunutsor S, Vitezova A, Oliver-Williams C, Chowdhury S, Kiefte-de-Jong JC, Khan H, Baena CP, Prabhakaran D, Hoshen MB, Feldman BS, Pan A, et al. Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *BMJ.* 2014; 348: g1903.
7. Bjelakovic G, Gluud LL, Nikolova D, Whitfield K, Wetterslev J, Simonetti RG, Bjelakovic M and Gluud C. Vitamin D supplementation for prevention of mortality in adults. *Cochrane Database Syst Rev.* 2014; 1: CD007470.